

Chapter 14

PROGNOSIS FOR EMPLOYMENT

The fourth question that we had initially asked ourselves was, Will the patient be able to earn a living for himself or will he remain dependent upon others?

The second Michigan Epilepsy Center follow-up project provided preliminary information, in this regard, from forty-six patients. The other patients involved in the project were either still in some type of school situation or were housewives. Twenty patients were employed at time of follow-up, and twenty-six were unemployed. The statistically significant correlates, with findings obtained at the time of the initial examination in regard to employment state at follow-up, are shown in Table 110. The

TABLE 110
SIGNIFICANT CORRELATIONS OF FINDINGS ON INITIAL EXAMINATION
WITH UNEMPLOYMENT AT TIME OF FOLLOW-UP

	<i>r</i>	<i>Significance Level (%)</i>
Attended special school	-.388	2
Full Scale IQ	-.366	2
Average marks in school	-.359	2
Amount of theta activity in EEG	.341	5
Performance IQ	-.339	5
Verbal IQ	-.332	5
"Organic" findings on Bender-Gestalt Test	.331	5
Prognosis for intellectual functions	.315	5

Seizure activity in EEG	.310	10
Generalized paroxysmal activity in EEG	.308	10
Focal grand mul seizures	.292	10
Prognosis for behavior	.281	10

significant correlates between findings obtained at follow-up examination and employment state are listed in Table 111. The coding system had been constructed in such a manner that "presently employed" was marked as 1 and "unemployed" as 2; the high end of the scale therefore represents unemployment. It is apparent that unemployment was mostly related to lower intellect and/or organic mental changes on the initial as well as follow-up examination. There is also a significant correlation

TABLE 111
SIGNIFICANT CORRELATIONS OF FINDINGS ON FOLLOW-UP EXAMINATION
WITH UNEMPLOYMENT AT TIME OF FOLLOW-UP

	<i>r</i>	Significance Level (%)
Organic mental changes	.477	1
Response to anticonvulsant medication	— .441	5
Serial 7 subtractions impaired	.415	1
Proverb interpretations concrete	.393	5
Recent memory impaired	.385	2
History of difficulty concentrating	.384	2
Behavior problems	.383	2
Verbal IQ	— .355	5
Full Scale IQ	— .339	5
Age	— .335	5
"Organic" findings on Bender-Gestalt test	.330	5
Remote memory impaired	.313	5
<hr/>		
Amount of theta activity in EEG	.289	10
Nonparoxysmal diffuse hyperventilation buildup in EEG	.253	10

with behavioral difficulties. While this may not be surprising, it is, of course, of considerable interest that variables related to seizure types or seizure frequency did not show significant relationships. The only hint that seizures may interfere with employment is contained in the correlate which shows poor response to anticonvulsant medication. Unemployed patients, therefore, are not likely to be in complete remission. Analysis of variance between the employed and unemployed groups, utilizing the previously mentioned 190 variables, confirmed—as the asterisks show—the findings of the correlations matrix and placed them

in somewhat sharper focus. Behavioral problems head the list, these were probably of long standing, as we noted in the previous chapter, and so were intellectual difficulties. They manifested themselves probably during school age in the form of poor grades and also necessitated special education in approximately half of the unemployed patients. The six-year difference in mean age between the groups is important because it suggests that some of the patients may be getting a slow start on the employ-

TABLE 112
SIGNIFICANT DIFFERENCES BETWEEN EMPLOYED AND UNEMPLOYED GROUP

	<i>Employed</i>	<i>Unemployed</i>	<i>P</i>	<i>Significance Level (%)</i>
Behavior problem at follow-up*	1.9	3.4	7.5	1
History of school grades*	4.6	3.2	6.1	5
Age	26.5 yrs.	20.6 yrs.	6.1	5
"Organic" findings on Bender-Gestalt test*	2.5	4.3	5.1	5
Performance IQ*	101.9	88.3	5.1	5
Prognosis for intellectual functions*	3.6	5.1	4.8	5
Amount of theta activity in EEG*	3.6	4.8	4.8	5
Full Scale IQ*	101.3	88.5	4.6	5
Digit Symbol—Wechsler IQ	8.6	6.8	4.4	5
Paroxysmal activity in EEG	1.1	2.3	4.4	5
Clusters of minor seizures for several days, freedom from seizures for several weeks	2.7	1.3	4.2	5

ment market; but as they get older and their seizure disorder improves somewhat, they begin finding jobs. The hard-core unemployed epileptic individual is, in all probability, the patient with lower IQ, and/or organic mental changes with or without associated personality difficulty. The fact that seizures do not necessarily preclude employment is demonstrated by the observation that the variable "clusters of minor seizures for several days with freedom from seizures for several weeks" occurred significantly more frequently in the employed than in the unemployed group in this particular sample of patients. As had been pointed out in the chapter on seizure prognosis, patients who

TABLE 113
SIGNIFICANT DIFFERENCES BETWEEN EMPLOYED AND UNEMPLOYED GROUP

		<i>Employed</i>	<i>Unemployed</i>	<i>X²</i>	<i>Signif- icance Level (C_t)</i>
Postictal headaches	Absent	3	12	6.2	5
	Present	16	8		
Attended special school*	Yes	3	14	5.2	5
	No	16	12		
Majority of tests were not applicable due to small numbers in sample.					
Rotation of Bender designs	Absent	16	13	3.6	10
	Present	3	12		
Academic difficulties in school	Absent	15	11	3.6	10
	Present	5	15		
Postictal confusion (major seizures)	Absent	17	12	3.0	10
	Present	2	8		
Postictal fatigue (major seizures)	Absent	7	14	3.0	10
	Present	12	6		
Focal major seizures	Absent	8	18	2.8	10
	Present	12	8		

present this phenomenon are not likely to achieve a permanent remission. It also deserves to be repeated that the variable "present seizure state" did not show significant differences between the two groups.

It has been mentioned previously that the Michigan Epilepsy Center had received a grant from the U.S. Public Health Service, Division of Vocational Rehabilitation, to study the aspects that relate to employment problems of epileptic patients. The material from that study will not be presented here because it merits extensive discussion which is beyond the scope of this publication. It did, however, provide an opportunity to check the main findings that were mentioned here. In the material from the VRA project there was likewise no significant differ-

ence in regard to seizure frequency between employed and unemployed patients. The detailed breakdown for the VRA patients is shown in Table 114. Two important aspects emerge: (1) The patient who has seizures less than once a year is, for practical purposes, in remission and is employed in the majority of cases. This is of course what one would expect on general grounds and (2) the more interesting point is that on the bottom of the scale there is no difference between the groups. A considerable number of epileptic patients, therefore, are able to maintain employment in spite of having, on the average, several

TABLE 114
FREQUENCY OF OCCURRENCE OF SEIZURES AT PRESENT IN RELATION
TO EMPLOYMENT STATUS

<i>Code Number and Description</i>	<i>Employed</i>	<i>Unemployed</i>
1 Less than once a year	32	12
2 About once a year	6	5
3 Two to three seizures a year	5	13
4 Four to six seizures a year	7	7
5 Seven to 12 seizures a year	4	6
6 Once a month	9	6
7 Two to three a month	11	9
8 Once a week	1	7
9 Several a week	13	10
<i>Totals</i>	88	84

seizures per week. The VRA study confirmed that seizures by themselves do not have to lead to unemployment. The major handicapping factors for the VRA group were likewise intellectual difficulties, organic mental changes, personality problems, and factors related to motivation.

The physician quite commonly hears the patient's complaint that he was fired because he had a seizure on the job, or he cannot find employment because "nobody will hire an epileptic." Although this may be the case in some instances, it has been my experience in virtually all cases where this was complained of by the patient, that subsequent mental status examination showed deficits which rendered the patient, for practical purposes, unemployable in a competitive work situation. The patient

may not be aware of these deficits, and it is much easier for him to blame his seizures for rejection by society, rather than organic mental changes, intellectual difficulties, or personality problems.

When the material from the VRA project is completely worked up, we should be able to provide formulas which will predict with reasonable accuracy the patient who is, in fact, unemployable in spite of all rehabilitative measures. More importantly, it should give the characteristics of those patients who are potentially employable, but for some reason or another have not succeeded in getting or maintaining a job. This would be the group towards which maximal rehabilitative efforts should be devoted.

Another interesting aspect of the findings is that they take the employment problems of the epileptic patient out of the strict realm of epilepsy (i.e. recurrent seizures) and place them in a more general framework. The factors that were found to be operative in the unemployed epileptic may well play a significant role in other hard-core unemployed individuals who tend to blame external circumstances rather than their own insufficiencies for their failure. It has been stated in the introduction to this book that the statistical methods used in this study to clarify aspects related to epilepsy could be applied without difficulty to any other condition one may want to investigate. Chronic unemployment is, in all probability, not only a sociological but, perhaps even more importantly, a medical problem in respect to the dimensions outlined above.

Chapter 15

INSTITUTIONALIZATION

A further question raised initially was in regard to the characteristics of patients who are likely to become permanent inmates of an institution for epileptics. The first follow-up study contained only five patients who had been institutionalized and the second study, seven patients. These numbers were too small for statistical purposes. We had, therefore, over the years transferred patients from Caro State Hospital for Epileptics to the Lafayette Clinic in order to get a clearer view of the problem. As previously mentioned in the chapter on seizure prognosis, fifty-seven of the Caro patients were included in the Lafayette Clinic inpatient review. These fifty-seven patients were contrasted against the 162 patients who had been referred from the community. Fortunately it turned out that there were no age or sex differences between the groups. The mean age was 26.6 years for patients from the community and 28.8 years for the Caro patients. There were eighty-six males and seventy-six females in Group I (patients from the community), and twenty-nine males and twenty-eight females in Group II (Caro State Hospital patients). Tables 115 and 116 show the variables that differed significantly between the two groups. The variables which are marked by an asterisk also showed significant differences in Tables 68 and 69 dealing with response to treatment in the hospital. The table containing the F tests demonstrates two main findings: (1) The Caro patients had an earlier onset and more intense seizure disorders than the patients referred from the community. There was a longer duration of illness, more frequent injuries during seizures, more different seizure types in

TABLE 115
SIGNIFICANT DIFFERENCES BETWEEN INSTITUTIONALIZED GROUP
AND PATIENTS REFERRED FROM THE COMMUNITY

	<i>Patients from Community</i>	<i>Patients from Caro State Hospital</i>	<i>F</i>	<i>Signif- icance Level (%)</i>
Duration of illness in years	12.2	22.7	45.8	1
Duration of main seizure type*	7.1	8.6	41.7	1
Amount of schooling*	3.8	2.4	41.0	1
Full Scale IQ*	83.6	64.0	36.8	1
Findings of cerebral pathology on neurological examination*	1.5	2.3	31.6	1
Frequency of injuries during seizures*	1.4	2.2	30.0	1
Focal atrophy on pneumoencephalography on the right	1.0	1.7	23.2	1
Age at time of onset of recurrent seizures in months*	170.7	84.1	21.7	1
Amount of alpha rhythm in EEG	2.8	2.0	19.9	1
Age at onset of first seizure in months*	151.6	65.3	19.5	1
Organic pathology suspected from psychological tests	2.5	3.1	17.3	1
Combination of seizures*	2.5	3.4	13.8	1
Evidence of bilateral cerebral disease*	1.9	2.3	12.8	1
Prenatal or perinatal injury suspected	1.4	1.9	12.5	1
Frequency of aura	2.6	1.8	10.7	1
Length of hospitalization in weeks*	6.8	9.2	8.8	1
Amount of theta activity in EEG*	2.8	3.3	8.5	1
Amplitude of background activity in EEG*	3.9	3.3	7.0	1
Amount of delta activity in EEG	1.1	1.3	6.9	1
Cerebral infection as etiological component	1.2	1.5	6.0	5
Frequency of maximal occurrence of seizures*	7.7	8.4	6.0	5
Neurotic tendencies on psychological testing	2.3	1.8	5.5	5
Clusters of seizures for several days, freedom from seizures for several weeks*	1.5	1.2	5.3	5
Amount of EEG abnormality*	3.4	3.8	4.9	5
Megimide induced seizure patterns, difficult to classify	2.8	3.7	4.8	5
Amount of fast activity in EEG	2.4	2.8	4.2	5
Evidence of left-sided cerebral disturbance from neurological examination and seizure patterns	1.4	1.8	4.0	5
Number of major seizures in hospital	3.1	6.6	6.4	5

TABLE 116
SIGNIFICANT DIFFERENCES BETWEEN INSTITUTIONALIZED GROUP
AND PATIENTS REFERRED FROM THE COMMUNITY

		<i>Patients from Community</i>	<i>Patients from Caro State Hospital</i>	X^2	<i>Signif- icance Level (%)</i>
Minor nonfocal motor seizures*					
	Absent	155	44	15.2	1
	Present	7	13		
Received regular schooling					
	Absent	9	14	14.7	1
	Present	144	39		
Focal grand mal seizures					
	Absent	92	19	9.2	1
	Present	70	38		
Theta rhythm present when patient has eyes opened*					
	Absent	151	46	5.9	1
	Present	11	11		
Focal grand mal variant seizure induced with Megimide*					
	Absent	121	37	5.4	5
	Present	8	9		
Received special schooling*					
	Absent	105	27	5.3	5
	Present	48	26		

the same individual, greater maximal frequency of seizures and more major seizures at the Lafayette Clinic. They also had more evidence of focal as well as diffuse cerebral pathology on clinical examination, pneumoencephalography, and psychological testing. The EEG mostly reflected cerebral damage with little alpha rhythm, more theta, delta and fast activity, and a greater amount of overall EEG abnormality. Amount of seizure patterns in the EEG, however, was not different between the groups, and (2) there were differences in regard to etiological factors. Prenatal and perinatal injuries, as well as cerebral infections, were significantly more common in the Caro group than in the group living in the community. A significant proportion of patients had been committed to the institution because of severe brain injury, making them unfit for life in the community, with epilepsy being superimposed.

We have repeatedly made the point before, that etiological factors did not seem to be of importance in regard to seizure control. The Caro State Hospital sample, however, showed that institutionalized patients had severe seizure disorders, and also more severe injuries to the central nervous system. The question arose, therefore, whether or not a greater intensity of the seizure disorder was a direct result of the more severe cerebral insults. To answer this question it was decided to divide the Caro State Hospital group into those patients in whom etiological factors had been present and into a group where no etiology could be demonstrated, and to compare the two groups on all variables. It should be mentioned at this time that the histories from Caro State Hospital were of excellent quality, containing very detailed information. In addition, we had in nearly all cases the opportunity to interview the parents after the patients had been transferred to the Lafayette Clinic. It was found that there were only twelve patients in whom absolutely no etiological factors could be implicated. When patients were included in whom one of the etiologies (prenatal or perinatal injury, postnatal head injury, cerebral infection, other significant external cause—e.g. cerebral malformation—, or family history of epilepsy) had been rated as questionably present, two groups could be formed. The first consisted of twenty-eight patients with no or questionable etiology; the other of twenty-seven patients with etiological factors. F tests and Chi Square tests were performed on the previously mentioned 190 variables. It was found that *no* statistically significant differences could be demonstrated between these two groups—apart from the variables dealing with etiology which had, of course, formed the basis of the separation into groups—except for the finding that a left-sided cerebral disturbance was found more commonly in the group in whom etiological factors were present (F 13.0, 1%). No significant differences were observed in regard to age at onset of illness, duration of illness, or to frequency, intensity or type of seizures. The IQ also was not significantly different (Full Scale 64 versus 61). This completely negative result was somewhat surprising and raised the question of whether we had contaminated our groups by including the cases of questionable etiology into those

with no discernible etiology, and by including family history of epilepsy as an etiological factor. It was therefore decided to take the twelve patients in whom no etiology had been present and contrast them with twelve patients who had had either definite cerebral infection or definite prenatal and/or perinatal injury. The major findings are shown in Table 117. Although we are dealing with a small sample, the trends are obvious. The patients with cerebral infection or perinatal injury were younger at the time of onset of their illness and younger in age at time of transfer to the Lafayette Clinic. This kept the duration of the illness essentially constant between the two groups. The group with known etiology also had an eleven-point lower mean IQ, but the intensity of the seizure disorder as expressed by remissions, frequency of maximal occurrence of seizures, frequency of seizures just prior to transfer to the Lafayette Clinic, frequency of seizures at the Lafayette Clinic, clusters of seizures in a given day, and occurrence of status epilepticus showed no differences between the groups. If there is a trend towards a difference at all, it would be that the group who had a definite etiology was actually slightly better off in respect to their seizure disorder. It is recognized that we are dealing with a very small sample of patients; nevertheless, there is no evidence that the intensity of the seizure disorder is related to the severity of cerebral injury.

Reviewing the patient material from Caro State Hospital, we noted that one is dealing essentially with three types of patients: (1) those who have an exceedingly intense seizure disorder, usually starting early in life, which does not yield to anticonvulsant treatment; (2) patients who have cerebral injuries, also usually acquired at an early age, leading to marked intellectual difficulties making them unfit for life in the community, and (3) a few patients with epilepsy whose main difficulty is not the intensity of the seizure disorder, or the severity of cerebral injury, but who show severe behavioral problems which cannot be tolerated by the environment.

The first group is the most interesting from the point of view of the epileptologist because it points to the existence of what one might call "malignant epilepsy." Fortunately this group is relatively small in size.

TABLE 117
COMPARISONS OF INSTITUTIONALIZED PATIENTS REGARDING THE INFLUENCE
OF ETIOLOGICAL FACTORS

	<i>No Etiology</i> (N = 12)	<i>Definite Etiology</i> (N = 12)
Mean age at time of first seizure	7.8 years	3.1 years
Mean age at onset of recurrent seizures	10.4 years	5.9 years
Duration of illness	26.7 years	26.0 years
Number of patients who were in remission at time of transfer to Lafayette Clinic	0	2
Number of patients who had experienced remission in the past	2	2
Number of different seizure types in a given patient:		
One seizure type only	3	4
Two seizure types	5	4
Three or more seizure types	4	4
Main seizure types most difficult to treat:		
Grand mal focal	5	9
Grand mal nonfocal	3	1
Psychomotor seizures	4	1
Grand mal variant nonfocal	0	1
Grand mal variant focal	0	1
Number of patients whose seizures started during first year of life	2	5
Duration of seizure type most difficult to treat	more than ten years	more than ten years
Maximal frequency of occurrence of seizures	several per week	several per week
Frequency of seizures prior to transfer to Lafayette Clinic	2 to 3 a month	7 to 12 per year
Number of patients who had clusters of seizures in one day	9	10
Number of patients who had clusters of seizures over several days, with freedom from seizures for several weeks	3	1
Number of patients with status epilepticus	3	2
IQ	69.2	58.7
Number of major seizures during hospitalization at Lafayette Clinic	64	52
Number of minor seizures during hospitalization at Lafayette Clinic	66	30

Chapter 16

LIFE EXPECTANCY

In the second MEC follow-up study we had noted that out of 136 patients whose fate could be ascertained, thirteen (9.6%) had died. This is a rather high proportion if one considers the age group that one is dealing with. As had been stated in the chapter on seizure prognosis, the mean age of patients at the time of death was 18.5 years if one excluded the three patients who had suffered from brain tumors, and 24.5 years if they were included. These are, of course, unusually low ages and are probably, at least in part, due to the small sample. Nevertheless, the figures did agree with the literature that the average life span of patients with epilepsy is lower than that of the general population.

In order to get some more information on this topic, we had asked the Michigan Department of Health, Section of Vital Records, to send us a listing of all the patients whose death certificates had shown epilepsy, for the years 1960 through 1966. We also inquired about the total number of deaths, the mean age at time of death for all individuals, and the mean age at time of death of patients who had seizures. Table 118 lists the findings. Age at time of death was expressed by the median rather than the mean because of infant mortality and the open-end age beyond ninety years. The table indicates that the diagnosis of epilepsy had appeared on the death certificate in approximately 0.1 per cent of all cases. It also shows that the median age at time of death of the patients in whom epilepsy was noted on the death certificates was approximately two and one-half decades below that of the average resident of the State of Michigan.

TABLE 118

NUMBER AND MEDIAN AGE AT TIME OF DEATH DUE TO ALL CAUSES IN THE STATE
OF MICHIGAN COMPARED WITH PATIENTS WHERE EPILEPSY WAS LISTED
AS DIRECT OR CONTRIBUTING CAUSE OF DEATH*

<i>Year</i>	<i>Total Deaths</i>	<i>Median Age at Death</i>	<i>Epilepsy</i>	<i>Median Age at Death</i>
1960	67,912	68.3 years	81	35.2 years
1961	67,375	68.7 years	74	40.8 years
1962	70,049	69.1 years	95	41.8 years
1963	72,438	69.3 years	91	34.7 years
1964	72,129	69.2 years	97	36.5 years
1965	73,665	69.5 years	109	41.8 years
1966	74,596	69.5 years	119	43.5 years

* Courtesy Michigan Department of Public Health, Vital Records Section

It has been pointed out in the discussion of the literature that studies based on death certificates alone will give only general trends rather than completely accurate data. The problem is that the presence of epilepsy frequently is not reported on the death certificates. The total number of patients who have epilepsy undoubtedly exceeds that which is reported on the certificates. This is not merely an assumption, as I failed to find the name of one of my former epileptic patients on the list supplied by the Michigan Department of Health. When the specific death certificate was requested it was noted that the cause of death had been listed as drowning and the diagnosis of epilepsy had not appeared on the death certificate. The wording on the death certificates in regard to significant chronic illnesses is actually so ambiguous that there is no particular reason to include the diagnosis for the majority of cases. The most relevant section of the Michigan death certificate is represented in Figure 24. The emphasis is obviously on conditions that are regarded as having directly contributed to the death of the person. Unless the patient's death is actually witnessed, it is impossible to say whether a convulsive seizure had led to the patient's death or not. There is no space provided on the death certificates to list the chronic illnesses the patient may have suffered, regardless of whether or not they were considered by the physician as a contributory cause of death. These points are made here to demonstrate that

19. CAUSE OF DEATH Enter only one cause per line for (a), (b), and (c) * This does not mean the mode of dying, such as heart failure, asphyxia, etc. It means the disease, injury or complication which caused death.		MEDICAL CERTIFICATION I. DISEASE OR CONDITION DIRECTLY LEADING TO DEATH*(a) _____ ANTECEDENT CAUSES Morbid conditions, if any, giving DUE TO (b) _____ rise to the above cause (a) stating the underlying cause last. DUE TO (c) _____ II. OTHER SIGNIFICANT CONDITIONS Conditions contributing to the death but not related to the disease or condition causing death.		Interval Between Onset and Death
19d. DATE OF OPERATION		19e. MAJOR FINDINGS OF OPERATION		20. AUTOPSY? Yes <input type="checkbox"/> No <input type="checkbox"/>
21a. ACCIDENT (Specify) SUICIDE HOMICIDE	21b. PLACE OF INJURY (e.g., in or about home, farm, factory, street, office bldg., etc.)	21c. (CITY, VILLAGE, OR TOWNSHIP)	(COUNTY)	(STATE)
21d. TIME OF INJURY (Month) (Day) (Year) (Hour) m.	21e. INJURY OCCURRED While at <input type="checkbox"/> Not While <input type="checkbox"/> Work at Work	21f. HOW DID INJURY OCCUR?		
22. I hereby certify that I attended the deceased from _____, 19____, to _____, 19____, that I last saw the deceased alive on _____, 19____, and that death occurred at _____ m., from the causes and on the date stated above.				
23a. SIGNATURE (Degree or title)		23b. ADDRESS		23c. DATE SIGNED

FIGURE 24. Copy of medical section of death certificate.

the Vital Statistics Section of the Michigan Department of Health has, at present, no way of knowing what the median age at time of death is for all patients who have suffered from epilepsy. When data based on death certificates are published in the future, it would be advisable to describe the relevant sections of the death certificates so that the reviewer can form an opinion about the completeness of the sample.

The most reliable way of gathering information on this topic is, of course, through follow-up of large numbers of patients for a long period of time, as is currently being done in Denmark. Our results from the review of death certificates are given here to demonstrate methodological problems, and so that they can be compared with other data that are being collected through follow-up studies. In spite of the difficulties which one encounters in the interpretation of data from death certificates, it is interesting that the findings do agree with the current study by Henriksen *et al.*, who had noted that the majority of their patients had died between the ages of thirty and forty-five years.

Chapter 17

SUMMARY

SEIZURE PROGNOSIS

Review of the literature showed that there has been no substantial improvement in *long-term* remission rates of epileptic patients over the past sixty years. Statements that approximately 80 per cent of patients are controlled by anticonvulsant medications are usually due to short follow-up periods and the inclusion of "improved" patients in the "controlled" group. A direct relationship has been shown to exist between length of follow-up and the percentage of patients who are seizure free at any given time. In regard to characteristics of patients which are likely to influence prognosis, the literature showed consensus on the following features: Patients who are normal in all other respects, except for the seizure disorder, tend to have a good prognosis especially if they have only grand mal seizures. Patients whose EEG is normal or normalizes with treatment can likewise expect a good outcome. The prognosis becomes poor with long duration of illness, a combination of different seizure types, large total number of seizures, and early age of onset.

Opinions are somewhat divided in regard to the importance of frequency of occurrence of seizures, whether seizures occur in the waking or sleeping state, heredity or other etiological factors, and abnormal findings on neurological examination.

It is repeatedly pointed out in the literature that patients with psychomotor seizures have a poorer prognosis than patients who have grand mal seizures. It should be borne in mind, however, that psychomotor seizure patients usually also have additional grand mal seizures and, therefore, one tends to compare patients

who have more than one seizure type with those who have only one seizure type. This fact should be considered in future research dealing with psychomotor epilepsy. It is apparent that major seizures have always responded better to anticonvulsant treatment than minor attacks, and modern drugs have not changed this pattern appreciably. The electroencephalogram has been shown to relate only in part to seizure prognosis. The relationship tends to improve if one looks at specific features of the tracings rather than relying on a global rating of normal versus abnormal. It is important to recognize that focal seizure discharges in a child's electroencephalogram do not necessarily represent the fixed atrophic type of lesion that one is accustomed to find in the adult.

In regard to febrile convulsions, the literature review indicates that they can be roughly divided into two groups: (1) simple or "benign" febrile convulsions and (2) epilepsy triggered by fever. It is interesting to note that essentially all the criteria that are given for differentiating between these two conditions also have been listed as being of importance in distinguishing between cases of epilepsy with good or poor prognosis.

Pyknolepsy was extensively discussed and it was demonstrated that the condition originally described under this name does not represent a clinical entity, and it seems to serve no purpose to perpetuate the term.

The prognosis of petit mal is still somewhat controversial, but the majority of authors feel that the seizures gradually decrease in number and intensity during adolescence and adulthood so that they do not form a major handicap for the patient later on. Precise figures for actual complete remissions are not readily available because the follow-up examination would have to include EEG evaluations in all instances. Historical reports about cessation of petit mal are unreliable. Approximately one-third to one-half of all patients starting with petit mal develop grand mal seizures later on. There is suggestive evidence that subsequent development of grand mal can be prevented by combined prophylactic treatment. This aspect deserves more intensive investigation using matched groups.

Infantile spasms—hypsarhythmia represent the most severe

form of childhood epilepsy. The success and failure of steroid treatment was discussed and the point was made that immediate treatment seems to be imperative if one wants to avoid the profound mental decay accompanying the condition. It was suggested that infantile spasms—hypsarrhythmia be regarded by the pediatrician and general practitioner as a medical emergency requiring immediate intensive treatment by specialists in the hope that permanent and severe loss of intellectual functions can be prevented.

In regard to posttraumatic epilepsy, there is still some controversy whether the patient's prognosis for long-term seizure remission differs from that of seizures due to other etiologies. Part of the difficulty resides in a problem of definition, namely, Does a single seizure after trauma constitute "epilepsy" or is the presence of several recurring seizures required for such a diagnosis? There tends to be agreement that the incidence of posttraumatic seizures stands in direct relationship to the severity of the cerebral injury. Penetrating wounds in the central-parietal regions appear to be most epileptogenic. The maximum incidence of posttraumatic epilepsy after wounds of this type was given in the literature as 65 per cent. This still leaves approximately one-third of patients who do not develop seizures after such injury. It seems therefore that cerebral damage, although a potent epileptogenic factor, cannot be sufficient cause for the development of a chronic seizure disorder. It is important to point out also that the probability for terminal remissions was shown, in two separate studies, to bear no relationship to the intensity of the injury. In general, the course of true posttraumatic epilepsy appears to be relatively mild with seizures spaced by fairly long intervals.

In contrast to medical treatment there has been definite progress in regard to surgical results. This is mostly attributable to the introduction of electroencephalography which demonstrated that patients with psychomotor seizures may have a single focal lesion in one temporal lobe. The surgical results do not seem to deteriorate in relation to length of follow-up as is the case with medical treatment. Operative intervention discloses, in a number of instances, previously unsuspected small but gross le-

sions which can be removed *in toto* with subsequent arrest of seizures. Unfortunately most patients with psychomotor seizures have multiple areas of pathological electrical activity, and only a relatively small number of patients become eligible for operation. Patients with focal discharges in areas other than the temporal regions tend to have a poorer surgical result. In severely disabled patients with infantile hemiplegias, uncontrolled seizures, and socially intolerable behavior, hemispherectomy appears to be of value. It was pointed out that neurosurgical follow-up criteria of patients have frequently been quite loose, and specific suggestions for improvements were made.

In order to determine long-term prognosis of seizure patients and the factors that are responsible for a good or poor outcome, two follow-up studies were conducted at the Michigan Epilepsy Center. The first was in the nature of a pilot study testing the data collection and data analysis procedures. Thirty-two children were involved who had been followed for at least five years after initial evaluation. A two-year terminal remission had occurred in 33.3 per cent, and a five-year terminal remission in 29.6 per cent. The second study dealt with ninety patients of all ages who had been followed for at least five years after initial evaluation. The two-year terminal remission rate was 32.2 per cent, and the five-year terminal remission rate was 16.7 per cent. These findings agree with the majority of studies published in the literature.

In order to arrive at prognostic indices for good versus poor outcome, the clinical and electroencephalographic findings were coded, punched on IBM cards, and subjected to the following statistical procedures: intercorrelation of findings, factor analysis, analysis of variance, and discriminant function analysis. Factor analysis produced four factors that were relevant for prognosis:

1. Epilepsy associated with cerebral damage.
2. EEG seizure activity.
3. Mixed psychomotor epilepsy.
4. Specific seizure propensity.

The following variables showed loadings on the various factors:

Ad 1. Patients with lowered IQ, "organic" features on psychological tests, abnormal findings on neurological examination, and slow landmarks of development showed

persistent seizures on follow-up and were unemployed.

Ad 2. The EEG was found to behave for the most part as an independent variable but was not likely to become normal if it showed seizure patterns at time of initial evaluation while the patient was already on some anticonvulsant medication. The factor loading in regard to seizure outcome was quite low.

Ad 3. Patients with psychomotor seizures in combination with other seizure types, in the presence of associated psychiatric difficulties, were found to have done poorly in regard to seizure outcome, behavior, and employment.

Ad 4. This factor dealt only with frequency of occurrence of seizures. This was shown to be unrelated to all other variables that were sampled, and this factor constitutes the core of the epilepsy problem.

Analysis of variance confirmed the findings obtained from the intercorrelation matrix. The most important variables that distinguished patients who had shown a two-year terminal remission from those who continued to have seizures were frequency of injuries during major seizures, number of different seizure types in the same patient, duration of seizure disorder, presence or absence of psychomotor seizures, presence or absence of clusters of seizures over several days followed by freedom from seizures for several weeks, degree of presence of seizure patterns in the initial EEG, and amount of overall EEG abnormalities.

Inasmuch as the patients in these samples had for the most part been treated by their family physicians rather than specialists in neurology, an attempt was made to compare these treatment results against findings obtained by the Neurology Outpatient Service of the Lafayette Clinic. One hundred twenty-three patients had been followed regularly for periods ranging between two and seven years. Seventy-four per cent of these patients had had a seizure within six months prior to their last clinic visit. In a subgroup of fifty-six patients who had been followed regularly for at least five years, a terminal remission of at least two years prior to last clinic visit had occurred in only 14.3 per cent of cases. It should be borne in mind, however, that the Lafayette Clinic Outpatient Service tends to see the more

difficult treatment problems, and patients who continue to attend outpatient clinics are likely to represent a negative selection. Findings from seizure clinics merely point to the existence of a group of hard-core, difficult to treat patients, but give no definite indication about the relative size of this group in comparison to the total epilepsy population.

Inasmuch as outpatients may or may not adhere to their prescribed drug regime, they do not readily lend themselves to a description of the characteristics of the patient whose seizure disorder is indeed resistant to anticonvulsant drugs. For these reasons, the results of treatment of 245 epileptic inpatients who had been hospitalized for at least three weeks on the Neurology Inpatient Service of the Lafayette Clinic were evaluated. One hundred thirty-two of these patients had been referred from the community because of uncontrolled seizures; eighty-two had been referred from the state hospital system of the State of Michigan for research and/or teaching purposes, and thirty-one had been admitted from the community because of marked behavioral difficulties. Of the 132 patients who had been referred from the community because of inadequate seizure control, 45.5 per cent did not have any seizures during hospitalization, 25.7 per cent had between one and three seizures, while 28.8 per cent had more than four seizures in spite of maximum treatment efforts. The subgroup of 132 patients, as well as the total group, were subjected to analysis of variance in order to delineate the characteristics of the patient who is unresponsive to maximum treatment efforts in the hospital. It was found that intensity of seizure disorder was the most important aspect in regard to treatment response. Presence or absence of presumed etiological factors was not of importance. The seven criteria that had differentiated between the remitted and the unremitted group in the long-term follow-up study of the Michigan Epilepsy Center also were found to have been important in the success or failure of short-term inpatient treatment. In addition, frequency of seizures immediately prior to hospitalization was important in regard to seizure control while hospitalized.

Discriminant function analysis based on these eight variables was performed. It classified correctly 76 per cent of the patients

who did not have seizures in the hospital and 81 per cent of those who had at least one seizure while on adequate medication. Cross-validation utilizing the sample of patients from the second follow-up study of the Michigan Epilepsy Center confirmed the results. The formulas for classifying the patients in regard to their likelihood of achieving seizure control, either in the hospital situation or as outpatients, were presented.

It was observed that the "seizure propensity" of the epileptic patient (i.e. tendency towards spontaneously recurring seizures) was not related to his "seizure threshold" as measured by the amount of Megimide needed to induce a clinical seizure. The implications of this finding were discussed from a theoretical as well as practical point of view.

Suggestions for the use of the discriminant function weights were made, especially in regard to selection of patients for inpatient treatment, and whether or not a patient's report about his current seizure frequency can be trusted. The weights were developed on predominantly adolescent and adult patients who already had been on some drug regime prior to their evaluation by a neurologist. Therefore, they will not necessarily be applicable to children or patients who have just experienced their first seizure.

The finding reported in the literature, that duration of seizure disorder prior to initial evaluation is important for prognosis, was confirmed. Eighty-five per cent of the patients seen in the first year had achieved a terminal remission of at least two years. If the patients were seen within the second year after the onset of their illness, the percentage of two-year terminal remissions had already dropped to 50 per cent, and if the illness had persisted for ten years or longer, the remission rate had dropped to 13 per cent. The possible implications for the pathophysiology of epilepsy were discussed and suggestions to study the phenomenon more intensively were made.

BEHAVIOR PROGNOSIS

As far as behavior is concerned, it was found in both follow-up studies that the tendency was towards improvement of ab-

normalities with the passage of time. Behavioral disturbances did not appear for the first time, in these samples, years after the seizure disorder had manifested itself; they either preceded the seizures or occurred at about the same time as seizures made their first appearance. Behavioral difficulties were most common and most persistent in the group of patients who had evidence of some form of cerebral damage. A discriminant function analysis was performed: 77.3 per cent of patients who did not have behavioral difficulties and 85.0 per cent of those who did were correctly identified. The variables related to the classification were Performance IQ, "organic" features on psychological tests, presence of seizures during the first year of life, history of behavioral difficulties in school, feeding problems in infancy, talking age, special schooling, average school grades, adverse social environmental factors, and objective findings of cerebral pathology on neurological examination. This aspect of the study represents preliminary data because it has not yet been cross-validated on another sample.

PROGNOSIS FOR INTELLECTUAL FUNCTIONS

The literature shows considerable controversy in this respect. There are only two points of definite agreement:

1. Epileptic patients can be subdivided into two groups. One might be called "epilepsy only," and the other, "epilepsy associated with brain damage of varying degrees." This latter group has definitely lowered intelligence as one might expect.
2. Epileptic patients show more fluctuation in upward or downward direction on test-retest measures than the normal population.

A further point upon which most authors are agreed is that patients whose seizures start in childhood have lower intelligence than those in whom the disorder manifests itself first in adolescence or adulthood. On the other points listed below there tends to be agreement between authors, but opinions to the contrary exist.

1. In cases of "epilepsy only," the intelligence level is

normalities with the passage of time. Behavioral disturbances did not appear for the first time, in these samples, years after the seizure disorder had manifested itself; they either preceded the seizures or occurred at about the same time as seizures made their first appearance. Behavioral difficulties were most common and most persistent in the group of patients who had evidence of some form of cerebral damage. A discriminant function analysis was performed: 77.3 per cent of patients who did not have behavioral difficulties and 85.0 per cent of those who did were correctly identified. The variables related to the classification were Performance IQ, "organic" features on psychological tests, presence of seizures during the first year of life, history of behavioral difficulties in school, feeding problems in infancy, talking age, special schooling, average school grades, adverse social environmental factors, and objective findings of cerebral pathology on neurological examination. This aspect of the study represents preliminary data because it has not yet been cross-validated on another sample.

PROGNOSIS FOR INTELLECTUAL FUNCTIONS

The literature shows considerable controversy in this respect. There are only two points of definite agreement:

1. Epileptic patients can be subdivided into two groups. One might be called "epilepsy only," and the other, "epilepsy associated with brain damage of varying degrees." This latter group has definitely lowered intelligence as one might expect.
2. Epileptic patients show more fluctuation in upward or downward direction on test-retest measures than the normal population.

A further point upon which most authors are agreed is that patients whose seizures start in childhood have lower intelligence than those in whom the disorder manifests itself first in adolescence or adulthood. On the other points listed below there tends to be agreement between authors, but opinions to the contrary exist.

1. In cases of "epilepsy only," the intelligence level is

within normal limits but shifted towards the lower end of the normal range.

2. Frequency of major seizures tends to be directly related to intellectual level.

3. A small but statistically significant decrease in intellectual functions can be seen on test-retest measures as far as group mean values are concerned.

4. The Performance areas of the Wechsler IQ scale appear to be more affected in epileptic patients than the Verbal areas.

The concept of "epileptic personality" was purposely omitted from consideration of the literature review because there exist only opinions and no reliably measured data.

The Michigan Epilepsy Center's second follow-up study contained fifty-six patients who had been tested twice on the Wechsler IQ scales with intervals ranging between five and nine years. On the initial tests, the mean Full Scale IQ was 94.5 with a Performance IQ of 95.7, and a Verbal IQ of 94.9. At the time of follow-up, the values were 93.6, 91.8, and 92.4. The decrease in Performance IQ was significant at the 1 per cent level of confidence, and that of the Verbal IQ at the 5 per cent level. A comparison of patients whose Full Scale IQ had decreased by seven or more points against those in whom the IQ had decreased less, remained stable or had increased, demonstrated that the patients with initially higher IQ levels tended to show decrease more commonly than patients in the lower intellectual range. This phenomenon could explain, in part, why there are proportionally fewer epileptic patients in the bright normal or superior range of intelligence than in the average population. The loss is usually not marked enough to place the patient into the dull-normal or borderline defective group. The point was made that a "normal" IQ score in the epileptic patient does not necessarily reflect his pre-illness intellectual level. The possible relations to emotional functions were discussed.

The Performance IQ was more affected by the seizure disorder than the Verbal IQ. It was demonstrated that seizures per se can lead to decrease in Performance IQ. The IQ tended to rise if the patient achieved a terminal remission of at least

two years. This was not the case if the seizure disorder was merely improved.

Discriminant function analysis was performed and criteria developed that would allow prediction as to whether a patient's IQ is likely to deteriorate. The formula classified correctly 95 per cent of patients whose IQ had dropped by seven or more points and 83 per cent of the others. The variables involved in the prediction were as follows: initial Full Scale IQ, number of different seizure types in the same patient, frequency of major seizures, clusters of seizures for several days—freedom from seizures for several weeks, "organic" features on psychological tests, presence of adverse social environmental factors, and EEG background amplitude. The formula has not been cross-validated as yet on another sample and will apply mostly to adolescents and adults rather than children.

Age at time of onset of the seizure disorder was not related to seizure outcome in this sample of patients. Definite relationships existed, however, between age of onset and intellectual functions. The mean Full Scale IQ of patients whose seizures started between birth and three years of age was 84.8, for the patients whose seizures started between four and twelve years it was 90.4, and for those who developed seizures between thirteen and twenty-seven years of age it was 101.3. A seizure disorder starting early in life, therefore, does not necessarily have a poor prognosis as far as seizure control is concerned but is likely to be associated with lowered intelligence, regardless of presumed etiology.

It is of considerable interest that modern statistical studies corroborated four of the points that were made by Turner in 1907 as being unfavorable in regard to intellectual functions:

1. Young age at time of onset of the illness.
2. Presence of major as well as minor seizures.
3. Frequent seizures.
4. Occurrence of the seizures in series.

EMPLOYMENT

Intercorrelation of findings on the patient sample from the second MEC follow-up study showed that employment problems

in the epileptic patients were related mainly to lower IQ, organic mental changes, and behavioral difficulties. It was surprising to note that seizure frequency was not related to the patient's employment state. This finding could be checked on another sample of 172 patients and was found to hold true, if patients whose seizure frequency was less than once a year were excluded.

INSTITUTIONALIZATION

The findings of fifty-seven patients from Caro State Hospital for Epileptics were contrasted against those of 162 patients referred to the Neurology Inpatient Service of the Lafayette Clinic from the community, in order to define the characteristics of patients who have to be institutionalized. It was found that the institutionalized patients tended to consist essentially of three groups: (1) patients with severe cerebral injuries, usually acquired early in childhood, (2) exceedingly intense seizure disorders, also usually starting early in life, and (3) some patients with epilepsy whose main difficulty is not the intensity of the seizure disorder or the severity of cerebral injury, but who show severe behavioral problems which cannot be tolerated by the community.

A comparison of institutionalized patients with known etiological factors against those in whom no recognizable etiology was present did not show appreciable differences between the groups. There was no evidence that the intensity of the seizure disorder was related to the intensity of cerebral injury.

LIFE EXPECTANCY

The literature review pointed out that the life expectancy of epileptic patients tends to be lower than that of the general population. Status epilepticus is still a significant cause of death, especially in patients who are institutionalized. There exists a group of patients with severe and uncontrollable seizures who die in early adulthood from generalized physical and mental deterioration. Death as a direct result from a seizure is quite

infrequent but does occur on occasion and is not preventable at all times. Inasmuch as patients with nocturnal seizures may roll over in bed and suffocate, the importance of supervised sleeping arrangements was stressed.

In order to check on the reports in the literature regarding life expectancy of epileptic patients, a survey of death certificates was carried out. The Michigan Department of Public Health calculated for us the annual median age at time of death for all persons who had died in the State of Michigan during the years 1960 through 1966, as well as that of persons where epilepsy was listed as primary or contributory cause of death on the death certificates. It was found that the median age at time of death from all causes ranged between 68.3 and 69.5 years. The median age at time of death for patients where epilepsy was mentioned on the death certificate ranged between 35.2 and 43.5 years. Surveys of this type have inherent methodological difficulties which were discussed and can only provide evidence for trends rather than leading to firm conclusions. Suggestions about improvements in the format of death certificates were made.

CONCLUSIONS

It has been mentioned in the Foreword that one of the main reasons for writing the book lay in the hope that some clues may be uncovered in regard to the pathophysiology of epilepsy. Reviewing the literature and my own investigations, it would seem clear that in the majority of instances epilepsy is *not merely* a symptom of a variety of other etiological conditions. By this I do not mean to deny the importance of the symptom of convulsive seizures in a patient who is developing a brain tumor, or in conditions such as tuberous sclerosis, Sturge-Weber syndrome, Unverricht's myoclonus epilepsy and the like. What is meant is that after a careful neurological evaluation has been performed, the majority of patients with chronic recurring seizures will not be found to have any other significant disease. Lest there be some misunderstanding on this point, I would like to reemphasize that a patient who starts with convulsive seizures at any age deserves the most careful neurological workup in order to eliminate the possibility of other neurological or metabolic illnesses; but the fact that convulsive seizures can be a symptom of a great variety of different illnesses affecting the central nervous system should not necessarily argue against the additional existence of a condition which I have called here "specific seizure propensity" of the individual. It is this aspect that deserves intensive study in the future. Although some type of injury to the central nervous system frequently can be demonstrated in patients with chronic seizure disorders, this is not a necessary nor sufficient cause for the disorder.

The difference between seizure threshold and propensity

towards spontaneously recurring seizures is likely to be of fundamental importance in the understanding of epilepsy. While a convulsive seizure is a symptom of a temporarily lowered threshold, it will remain isolated or infrequent unless the individual also has the necessary mechanisms for propensity towards spontaneously recurring attacks. The intensity of that latter factor is likely to make the difference between patients who have occasional seizures only and patients who are not controllable by current anticonvulsant treatment. It also may well make the difference between children who have what is now called "benign" febrile convulsions and "epilepsy triggered by fever."

Hughlings Jackson once pointed out that when confronted with an epileptic patient ". . . the first question in my mind is not 'Is it a case of epilepsy?' but 'Where is the lesion permitting occasional excessive discharge?'" This was indeed a giant step forward nearly one hundred years ago, but we should not continue merely to echo Jackson's thoughts. The great majority of neurophysiological and neurochemical investigations still deal with the "epileptogenic focus" or the properties of the "epileptogenic neuron." These are important studies, but they are likely to be insufficient in providing the final answer to the problem. In addition to Jackson's question, one should also ask one's self, What are the factors that are responsible for the spread of abnormal electrical activity in this particular patient? Even more important would be the question, How does the patient's condition differ on the five days of the week when he is seizure-free from that of the sixth day when he has an attack? The neurophysiological and neurochemical events that lead up to this sixth day of the overt seizure would deserve at least equal attention as that which is being paid to the epileptogenic focus. The focus, after all, tends to be a more constant event rather than an intermittent one. Studies which deal with the cerebral environment in which the epileptogenic focus finds itself on different days would certainly be most valuable and would deserve a high priority. It bears repeating that the main problem in chronic epilepsy is not only the epileptogenic focus or the paroxysmal discharge, but the abnormal spread of electrical activity to other parts of the brain and/or the periphery. The presence of a focus

or diffuse bursts tends to lower the seizure threshold, but those factors which allow for spread are the ones that are in all probability related to the seizure propensity of the individual. As a crude analogy, one might point to the development of a forest fire. A lighted cigarette carelessly thrown away in the woods during a hot and dry summer will result in a fire, the extent of which will depend on the preceding length of absence of rain. The same lighted cigarette thrown away in the same location during the rainy season will extinguish itself harmlessly. Let us therefore not remain obsessed only with the cigarette or focus, but consider in much greater detail the state of the rest of the cerebral forest. The investigations of focal epilepsy have brought about great improvements in our concepts about the disorder, but they remain incomplete. One has to see the entire spectrum of the condition, ranging from the mildest cases to severely demented individuals who still occupy state hospital beds, in order to fully appreciate the complexity of the issue.

The differentiation of epilepsy into *idiopathic* and *symptomatic* has not been very fruitful in the past and has not been shown to be of prognostic significance. One of the reasons for this is probably the fact that the terms do not designate homogeneous groups. The word "idiopathic" is frequently used synonymously with genetic and at other times it refers to patients in whom no etiology can be found by careful history. The fact that no etiology is discernible by history does not necessarily mean that the patient could not have a "small but gross" lesion in one temporal lobe, which is discovered only incidentally, as a result of temporal lobectomy for seizure control. The word "symptomatic" implies that the etiology lies in some form of injury to the central nervous system, yet in a number of instances this "injury" is only a relatively minor accident which may have nothing whatsoever to do with the appearance of the seizure disorder. In other instances a definite family history of epilepsy is available in clearly symptomatic cases where there also has been undeniable brain injury. It is an open question to what extent the two elements conspired to produce a seizure disorder which may or may not respond to treatment. While brain injury lowers seizure threshold, it has not been shown that it affects seizure

propensity. This would explain why the same amount of injury will produce only isolated seizures in one individual and a severe intractable disorder in another.

The "constitutional" factor in epilepsy has of course long been recognized but has recently been rather neglected. Although we have at present no way of knowing what this constitutional factor is, it serves no purpose to deny its existence because research efforts will then be thwarted. Part of the reason for omitting the constitutional element from consideration seems to have been that the word has been equated by some authors with "hereditary." This is unfortunate because it narrows the term and suggests a mechanism which so far has not been proven to be operative in the great majority of patients. There is definite evidence that certain EEG abnormalities are inherited in a dominant mode. There is also definite evidence that patients who have these EEG abnormalities are likely to have a lowered threshold for seizure induction, but—and this is the important point—there is no evidence that these patients also have a propensity towards spontaneously recurring seizures. Siblings of epileptic patients—especially of those who have spike wave activity in their EEGs—have a high incidence of EEG abnormalities but only a very low incidence of overt recurrent convulsive seizures. It is clear that some additional factor must be operative in the epileptic patient on which he differs from his siblings. If we accept the concept that the resting EEG is related to seizure threshold rather than seizure propensity, we could explain also the occasional paradox where under treatment the EEG actually may become worse but the patient better.

Classifications of epilepsy based on presumed pathophysiological mechanisms are not likely to be useful in the immediate future because of our ignorance in regard to the really important basic factors. Until the time comes that we do have the necessary understanding, we shall probably be well advised to stay on a purely descriptive level of the clinical condition. Although the term "epilepsy" has nothing to commend itself—except usage through the ages—it seems to serve little purpose to coin new words at this time which suggest knowledge that is not available.

A distinction between patients with "epilepsy only" and "epi-

lepsy associated with cerebral damage" of varying degrees may sound on the surface like a resurrection of the idiopathic versus symptomatic classification. This is, however, not the case. The term "epilepsy only" means that the patient is healthy in all other respects to the best of our knowledge. It does not imply the presence or absence of presumed etiological factors or any pathophysiological mechanism. The patient who is classified as having epilepsy associated with mild, moderate, or severe cerebral damage has obviously more than one handicap. The term does not indicate that cerebral damage caused the seizures; it could be a result of them, or it may actually be independent of the seizure disorder. Although a distinction of this type would not necessarily yield a prognosis for seizure cessation, it could be useful in forecasting the patient's overall life achievements.

It has been pointed out that the chances for a patient to achieve complete seizure freedom rest mainly on the intensity of his seizure disorder and its duration. There is possibly some relationship between these aspects. A severe seizure disorder tends to be refractory to treatment by the time the neurologist sees the patient. The question arises, however, whether the severity could have been influenced by more intensive treatment at the very onset of the illness. The finding that 85 per cent of patients who are seen by a specialist within the first year of their illness achieve a terminal remission of at least two years is most impressive. Even more important is the observation that the results drop to 50 per cent after the first year. Combining these findings with Lund's observations that treatment results of the first three months herald the future course of the disorder, a strong case could be made that a maximum effort should be exerted to control seizures at the very onset. The medication regime should be adequate for the age and weight of the patient and rearranged with each successive seizure. There would seem to be no room for complacency in this regard. Follow-up appointments should initially be closely spaced, and comparison EEGs should be obtained at each visit.

The problem is, however, that at the present time we have no way of knowing whether a seizure in a child or adult will remain isolated or whether it represents the beginning of a chronic sei-

zure disorder. A differentiation of these two conditions would certainly be most important and specific matched control studies dealing with the total evaluation of patients immediately after their first seizures are indicated. All of the currently used anti-convulsants tend to have some side effects in a significant number of patients when given in adequate dosages. Their indiscriminate use therefore is not to be advocated, but hesitation to use them is equally dangerous if they are indeed capable of preventing an individual from becoming a chronic epileptic patient. This is a dilemma the physician faces today, and it is regrettable that there are so far no rigorous studies available that could provide us with guidelines in this respect.

We must not be satisfied with treatment results that lead to "improvement" but not to cessation of seizures. The fact that intelligence tends to decrease if the illness persists unchecked has to be taken into account. This is even more important when we recognize that the patients with an initially bright normal or superior intelligence quotient are more likely to suffer loss of intellectual functions. A basically bright person confronted with the fact of decreasing intellectual functions cannot help becoming a problem to himself or to society. These aspects of the epilepsy issue tend to be glossed over at the present time when we hear of improvement in seizure control that is being achieved by modern drugs. We have to continue to search for better treatment, by drugs or other methods, that will lead to complete cessation of attacks. While intellectual decrease is a most serious event at any stage of life, it becomes a virtual catastrophe if it occurs in childhood. The physician has been trained to regard conditions as emergencies where the life of the patient hangs in the balance. We can now save lives but we are woefully inadequate in saving that which makes us specifically human—namely, the intellect. The saving of intellectual functions will have to assume greater urgency than it has been given in the past. While this applies to all patients, it could perhaps be most clearly seen in the cases of infantile spasms-hypsarhythmia. A delay of treatment of a few *days* might make the difference between a child who will become either an inmate of a state institution or a self-sustaining citizen. This condition would surely

qualify for emergency measures in spite of the fact that life itself is not at stake. Inasmuch as the specialist does not tend to see these children until later in the course of their illness, by which time irrevocable damage is likely to have occurred, it becomes imperative that general practitioners and pediatricians who represent the first line of medical defense become thoroughly familiar with the diagnosis and its potential implications. Treatment, complex as it is, may well be left in the hands of the specialist provided he is called to the scene immediately and can see the child within a matter of a few hours.

While these comments dealt with the duration of the illness and its relationship to treatment response, there also appears to be a relationship between age at time of onset and the severity of the disorder. We have not been able to demonstrate in our own studies that patients whose seizures start before the age of three years develop necessarily a chronic intractable disorder. This does not rule out, however, that a more indirect relationship might exist. It could be postulated that a severe seizure disorder is likely to manifest itself early in life, but not every instance of epilepsy starting in childhood has to be of this severity. The child, as a result of maturational factors, has *a priori* a lower seizure threshold than the adult. He may therefore respond with a seizure to less provocation than the older individual. A milder seizure tendency may bring on clinical seizures in the child which may promptly respond to anticonvulsant medication. The seizure tendency apparently decreases with increasing age, and this may well be the reason why patients with "late onset epilepsy" and posttraumatic epilepsy of adult life seem to have relatively infrequent seizures. This concept could explain why different authors come to different opinions about the prognostic importance of age at time of onset of the illness. It could also explain the observation of our own investigations that children are somewhat more likely to show either complete seizure cessation or chronic disorders rather than occupying a middle position of some improvement as time goes on.

With epilepsy being a condition of such diverse clinical manifestations and markedly varying intensity, it is apparent that any general rule about what epileptic patients can or cannot do is

bound to be unjust. Each patient deserves careful individual attention in regard to his assets as well as his liabilities. Restrictions that have to be applied should not hold across the board for all patients, but should deal only with a particular individual at a particular time. They should be reevaluated periodically in regard to their continued need, with the goal being maximal opportunity in the presence of minimal risk. In actual practice this means that some epileptic patients can be leading essentially normal lives, while others may not be able to operate motor vehicles and work around dangerous machinery, while still others may need more or less constant supervision by members of the family or in a hospital environment.

This book has emphasized, as mentioned in the Foreword, the areas where progress has been slower than one might have been led to believe. This was done in order to point out our deficiencies and pave the way for a more rapid pace of accomplishments, but it does present a potential danger. There has been over the past several decades a progressive liberalization of rules and laws in regard to epilepsy. Many dedicated persons have spent a great deal of effort to remove the stigma that is still attached to this diagnosis. These efforts are most important and are just beginning to bear fruit in a variety of ways. The findings reported in this book should not be used to undermine them. Baseless pessimism in regard to the condition is at least equally as dangerous as false optimism. As long as we are unable to cure the majority of our epileptic patients, we should at least try to make their lot as easy as humanly possible, while persisting in our efforts directed toward the permanent eradication of the disorder. At the present time, epilepsy still represents just as great a challenge as it has throughout the ages. By fully recognizing this challenge and bringing to bear on it, in addition to the most modern equipment, our best scientific minds, we should be able to find the way that will lead not only to the permanent cure of those patients who already suffer from the illness, but also to its prevention.

APPENDIX

VARIABLES USED IN FIRST MEC
FOLLOW-UP PROJECT CORRELATION MATRIX

INITIAL FINDINGS

Age
Sex
Physical Health
*Duration of Labor
*Forceps Delivery
*Birth Difficulty
Feeding Problems in Infancy
Colic in Infancy
Crying During First Year of Life
Patient's Weight Gain Progress in Infancy
*Activity During First Year of Life
Sitting Up Age
Walking Age
Talking Age
*Delayed Maturation
*Seizures Present During First Year of Life
*Highest Degree of Fever Attained
*Febrile Convulsions
Nonfebrile Convulsions in Infancy
*Bedwetting
Usual Childhood Diseases without Delirium
*Behavior Difficulties in School
*Academic Difficulties in School
Personality Disorder
Family History
Infantile convulsions without fever
*Epilepsy
*Diabetes

*Temporary psychiatric hospitalization

*Chronic alcoholism

Seizure Type

*Nonfocal major

*Focal major

*Minor focal motor

*Psychomotor

***Combination of Seizures**

***Prognosis for Seizure Control**

***Prognosis for Behavior Control**

***Prognosis for Academic Achievement**

Major Seizures

Duration since onset

Present since first year of life

*Frequency of occurrence

*Remission in the past

Clusters of seizures in one day

*Status epilepticus

Relationship of time of day to seizures

Occurring within two hours after awakening

Frequency of occurrence of aura

Loss of bladder control

*Duration of one attack

State of Consciousness During Seizure

Dazed

Completely unconscious

Not altered

Posture During Seizure

Falls suddenly

No movements observed

Patient stiff

Twitching or shaking

Facial Color During Seizure

Pale

Cyanotic

Prominent Symptoms of Seizure

Blank stare

Head turns to right
Swallowing motions
Vocalization
Right arm or hand twitching
Left face twitching
Left arm or hand twitching
Left leg twitching
Postictal State
Confusion
Sleep
Unilateral muscle weakness
Nausea
Minor Seizure
Duration since onset
Frequency of occurrence
Remission in the past
Clusters of seizures in one day
Status epilepticus
Clusters of seizures over several days, freedom from seizures
for several weeks
Relationship of time of day to seizures
Frequency of occurrence of aura
Loss of bladder control
Duration of one attack
Dazed
No aftereffects
***Social Factors Contributing to Illness**
Etiological Factors in Neurological History
***Objective Findings of Cerebral Pathology on Neurological Examination**
***Psychological Test Results**
Intellectual level
Immaturity
Neurotic tendencies
Psychotic tendencies
Personality disturbances
***Organic pathology suspected**

Age at Onset of Convulsive Disorder*Duration of Illness*****Highest Rise of GTT From Fasting****EEG*****Amount of abnormality*****Amount of seizure patterns*****Amount of focal abnormalities*****Amount of alpha activity*****Amount of theta activity*****Amount of fast activity****Generalized paroxysmal activity****Intervals between paroxysms****Amplitude of background rhythms, Left****Amplitude of background rhythms, Right****Main background frequency****Adequate Effort but No EEG Change with Hyperventilation****Final Diagnosis****Convulsive disorder****Psychiatric disorder in addition to diagnosis of epilepsy****Mental retardation in addition to diagnosis of epilepsy****FOLLOW-UP FINDINGS****EEG*****Amount of abnormality*****Amount of seizure patterns*****Amount of focal abnormalities*****Amount of alpha activity*****Amount of theta activity*****Amount of fast activity****Generalized paroxysmal activity****Intervals between paroxysms****Amplitude of background rhythms, Left****Amplitude of background rhythms, Right****Main background frequency****Adequate Effort but No EEG Change with Hyperventilation****Patient On or Off Anticonvulsant Medication**

- °Present Seizure State
- °Behavior Problem
- °Academic School Problem
 - Attending or Attended Regular School
 - Attending or Attended Special School
 - No Formal Education
- °Average Marks
 - Grade Failure
- °Institutionalized

°The variables marked by asterisk were used in the factor analysis.

VARIABLES USED IN SECOND MEC
FOLLOW-UP PROJECT CORRELATION MATRIX

INITIAL FINDINGS

- *Sex
 - Physical Health
 - Difficulty During Pregnancy of the Mother
 - Duration of Pregnancy
 - Duration of Labor
- *Birth Weight
- *Condition of Child at Birth
 - Birth Cry
 - Condition of Patient's Head at Birth
- *Birth Difficulty
 - Feeding Problems in Infancy
 - Colic in Infancy
 - Activity During First Year of Life
- *Seizures Present During First Year of Life
- *Sitting Up Age
 - Walking Age
- *Talking Age
 - Toilet Training
 - Dry at Night
 - Highest Degree of Fever Attained
- *Febrile Convulsions
 - Nonfebrile Convulsions in Infancy
- *Bedwetting
- *Behavior Difficulties in School
 - Amount of Schooling
- *Attending or Attended Special School
 - Average Grades

***History of Personal Relations During Adolescence**

Family History

- *Stillbirth
- *Early infantile deaths
- *Infantile convulsions with fever
- *Infantile convulsions without fever
- *Epilepsy
- *Diabetes
- *Chronic alcoholism

Seizure Type

- *Nonfocal major
- *Focal major
- *Minor focal motor
- *Psychomotor
- *Absence
- *Combination of Seizures

Etiology of Seizures

- *Unknown
- *Birth injury
- Postnatal head injury
- *Cerebral infection
- *Mixed hereditary and external cause
- *Hereditary cause only

Prognosis for Seizure Control

Prognosis for Behavior Control

Prognosis for Academic Achievement

Adequacy of Medication Regime in the Past

Never Treated Before

- *Patient's Initial Response to Adequate Amount of Anti-convulsants

***Duration of Seizure Disorder**

Major Seizure

- *Frequency of occurrence, Maximal
- *Frequency of occurrence at present
- *Remission in the past
- *Clusters of seizures in one day
- *Status epilepticus

Relation of menses to seizure

EEG

- *Amount of abnormality
- *Amount of seizure patterns
- *Amount of focal abnormalities
 - Amount of alpha activity
 - Amount of theta activity
 - Amount of fast activity
- Right temporal focus
- Left temporal focus
- Focus consists of theta discharge
- Focus consists of complex discharge
- *Focus consists of sharp wave or spike
- *Generalized paroxysmal activity
- *Amount of spike wave activity
- *Amplitude of background rhythms, Left
 - Main background frequency
- ***Social Factors Contributing to Illness**
- ***Objective Findings of Cerebral Pathology in Neurological Examination**

Psychological Test Results

- *Immaturity
- *Neurotic tendencies
- *Psychotic tendencies
- *Personality disturbances
- ***Final Diagnosis Psychiatric Disorder in Addition to Diagnosis of Epilepsy**
- ***Organic Brain Pathology Suspected on Bender-Gestalt Test**
 - Rotation of 45 Degrees or More on Bender-Gestalt Test
- ***Verbal IQ Score**
- ***Performance IQ Score**
- ***Full Scale IQ Score**
- ***Difference Between Verbal and Performance Score**

FOLLOW-UP FINDINGS

- *Age
 - Seizure Type

Nonfocal Major

Focal Major

Minor Focal Motor

Psychomotor

Absence

Combination of Seizures

°Adequacy of Medication Regime in the Past

Response to Adequate Amount of Anticonvulsant Medications

Change in Number of Seizure Types

Remission at Present for Major Seizures

°Remission in the Past for Major Seizures

Remission at Present for Minor Seizures

Remission in the Past for Minor Seizures

°Remote Memory

°Recent Memory

°Serial 7 Subtractions

°Proverb Interpretation

Egocentric or Bizarre Interpretation

°Organic Mental Changes

Difficulty Concentrating

°Personality Problem

Sociopathic Behavior and/or Antisocial Behavior in P.I.

Depression

Destructive or Assaultive Behavior

EEG

°Amount of abnormality

°Amount of seizure patterns

°Amount of focal abnormalities

°Amount of alpha activity

°Amount of theta activity

°Amount of fast activity

°Right temporal focus

°Left temporal focus

Focus consists of theta discharge

°Focus consists of complex discharge

°Focus consists of sharp wave or spike

°Generalized paroxysmal activity

- *Amount of spike wave activity
- *Amplitude of background rhythms, Left
 - Amount of nonparoxysmal buildup, HV
 - Amount of paroxysmal buildup, HV
- *Amount of photic driving at low flash rates
- *Amount of photic driving at medium flash rates
- *Amount of photic driving at high flash rates
 - Main background frequency
 - Abnormalities improved or deteriorated
- *Organic Brain Pathology Suspected on Bender-Gestalt Test
 - Rotation of 45 Degrees or More on Bender-Gestalt Test
- *Verbal IQ Score
- *Performance IQ Score
- *Full Scale IQ Score
- *Difference Between Verbal and Performance Score
 - Verbal Scale Score Difference Between Evaluation I and II:
 - Worse
 - Better
 - Verbal Scale Score Difference Between Evaluation I and II:
 - Worse
 - Better
 - Performance Scale Score Difference Between Evaluation I and II:
 - Worse
 - Better
 - Performance Scale Score Difference Between Evaluation I and II:
 - Worse
 - Better
- Interval Between Two Visits
- Handedness
- *Present Seizure State
 - Patient On or Off Anticonvulsant Medication
- *Behavior Problem
 - Academic School Problem
 - Attending or Attended Special School
 - Average Marks in Regular School
- *Gainfully Employed
 - Housewife Performing Duties Adequately
- *Overall Condition of Patient

*Variables carrying asterisk were used for factor analysis.

VARIABLES USED IN SECOND MEC
FOLLOW-UP PROJECT

F-TESTS

Age
Physical Health
Duration of Labor
Birth Weight
Condition of Child at Birth
Length of Time Till Mother Saw Child after Delivery
Feeding Problems in Infancy
Crying During First Year of Life
Activity During First Year of Life
Sitting Up Age
Walking Age
Talking Age
Toilet Training
Age at Which Dry at Night
Highest Degree of Fever Attained
Duration of Fever over 104 Degrees
Amount of Schooling
Average Grades
School Truancy
Personal Relationships During Adolescence
Etiology
Birth injury
Cerebral infection
Mixed hereditary and external cause
Hereditary cause only
Major Seizures
Duration since onset

Frequency of maximal occurrence

Frequency of occurrence at present

Remission in the past

Clusters of seizures in one day

Status epilepticus

Clusters of seizures over several days, freedom for several weeks

Relationship of time of day to seizures

Relation of menses to seizures

Frequency of occurrence of aura

Loss of bladder control

Tongue biting

Injuries sustained during attack

Duration of one attack

Minor Seizures

Duration since onset

Frequency of maximal occurrence

Frequency of occurrence at present

Remission in the past

Clusters of seizures in one day

Clusters of seizures over several days, freedom for several weeks

Relationship of time of day to seizures

Frequency of occurrence of aura

Duration of one attack

Combination of Seizures

Initial Response to Anticonvulsant Medications—

1-3 Months

Subsequent Response to Anticonvulsant Medications—

3 Months-1 Year

Subsequent Response to Anticonvulsant Medications—

1 Year and after

Seizure Pattern Change after Anticonvulsant Treatment EEG

Amount of abnormality, waking record

Amount of abnormality, waking and sleep record

Amount of seizure patterns

Amount of focal abnormalities
Amount of alpha activity
Amount of theta activity
Amount of fast activity
Amount of diffuse delta activity
Amount of paroxysmal activity
Left background rhythms amplitude
Right background rhythms amplitude
Main background frequency
Wechsler IQ
Verbal Scale
Performance Scale
Full Scale
Information
Comprehension
Arithmetic
Similarities
Digit Span
Vocabulary
Digit Symbol
Picture Completion
Picture Arrangement
Block Design
Object Assembly
Organic Brain Pathology Suspected from Bender-Gestalt Test
Forty-five Degree or More Rotation of Bender-Gestalt Figures
Psychological Tests
Immaturity
Neurotic tendencies
Psychotic tendencies
Personality disturbances
Organic pathology suspected
Social Factors Contributing to Illness
Laboratory Findings Summary
Objective Findings of Cerebral Pathology on Neurological Examination
Etiological Factors Summary

Prognosis for Seizure Control

Prognosis for Behavior Control

Prognosis for Academic Functions

Follow-Up Results

Present seizure state

Behavior problem

Academic school problem

Average marks

Overall condition of patient

VARIABLES FROM SECOND MEC FOLLOW-UP PROJECT

CHI SQUARE TESTS

Sex

Pregnancy Complications

Marked nausea and vomiting

Labor Complications

Forceps delivery

Birth cry, induced versus spontaneous

Colic in Infancy

Attended Special School

Seizures Present During First Year of Life

Birth Difficulty

Febrile Convulsions

Bedwetting

Behavior Difficulties in School

Academic Difficulties in School

Severe Infectious Disease

Severe Head Injury with Unconsciousness

Neurotic Symptoms

Personality Disorder

Family History

Stillbirths

Breech deliveries

Multiple births

Early infantile deaths

Congenital defects

Mental retardation

Temper tantrums

Behavior problems

Academic school problems
Meningitis
Bedwetting
Breathholding spells
Infantile convulsions with fever
Infantile convulsions without fever
Epilepsy
Fainting spells
Severe chronic headache
Chorea
Diabetes
Excessive nervousness
Nervous breakdown
Temporary psychiatric hospitalization
State hospitalization
Chronic alcoholism
Abortions or miscarriages

Etiology Unknown**Seizure Type**

Nonfocal major
Focal major
Minor focal motor
Psychomotor
Absence

Major Seizures

Present since first year of life
Occurring at night during sleep
Precipitating event—Nonspecific excitement
Precipitating event—Emotional stress
Precipitating event—Omission of anticonvulsant medications
Posture during attack—Slides gradually to ground
Posture during attack—Falls suddenly
Muscular action during attack—Classic tonic and clonic phase
Muscular action during attack—Patient stiff
Muscular action during attack—Twitching or shaking
Facial color during attack—Pale
Facial color during attack—Cyanotic

Eyes rolled up during attack

Postictal state—Fatigue

Postictal state—Headache

Postictal state—Confusion

Postictal state—Sleep

Postictal state—Nausea

Prominent feature of attack—Salivation

Prominent feature of attack—Vocalization

Prominent feature of attack—Right side twitching

Prominent feature of attack—Left side twitching

Minor Seizures

Occurring within 2 hours of awakening

Emotional stress as precipitating event

Consciousness during attack—Not altered

Consciousness during attack—Hears, but cannot respond

Consciousness during attack—Dazed

Consciousness during attack—Completely unconscious

Posture during attack—Not altered

Postictal state—No aftereffects

Postictal state—Sleep

Feature of attack—Random wandering around

Adequacy of Medication Regime in the Past

Reason for Referral—Emotional or behavior problems

Diagnosis of Psychiatric Disorder in Addition to

Diagnosis of Epilepsy

Psychiatric Treatment Recommended

EEG

Right temporal focus

Left temporal focus

Type of discharge—Sharp wave or spike

Forty-five Degree Rotation in Any Figure on Bender-Gestalt Test

Follow-Up Results

Takes anticonvulsant medication

Attending regular school

Attending special school

Grade failure

Gainfully employed

ESTIMATED PROBABILITY OF CORRECT CLASSIFICATION (2 GROUPS)

DIFFERENCES BETWEEN DISCRIMINANT FUNCTIONS	PROBABILITY
.00	.50
.04	.51
.08	.52
.12	.53
.16	.54
.20	.55
.24	.56
.28	.57
.32	.58
.36	.59
.40	.60
.45	.61
.49	.62
.53	.63
.57	.64
.62	.65
.66	.66
.71	.67
.75	.68
.80	.69
.85	.70
.89	.71
.94	.72
.99	.73
1.04	.74
1.10	.75
1.15	.76
1.21	.77
1.26	.78
1.32	.79
1.39	.80
1.45	.81
1.52	.82
1.58	.83

Classification Probability

371

1.66	.84
1.73	.85
1.81	.86
1.90	.87
1.99	.88
2.09	.89
2.20	.90
2.31	.91
2.44	.92
2.59	.93
2.75	.94
2.94	.95
3.18	.96
3.48	.97
3.89	.98
4.59	.99

CONSTANTS AND WEIGHTS FOR PREDICTING INPATIENT TREATMENT RESULTS

	DISCRIMINATE FUNCTION I								
	1	2	3	4	5	6	7	8	9
Constant	-17.43								
Injuries during seizures	-0.41	-0.82	-1.23	-1.64	-2.05	NA	NA	NA	NA
Combination of seizures	-0.39	-0.78	-1.17	-1.56	1.95	-2.34	-2.73	-3.12	-3.51
Psychomotor seizures	6.08	12.16	NA	NA	NA	NA	NA	NA	NA
Duration of seizure disorder	2.85	5.70	8.55	11.40	14.25	17.10	19.95	22.80	25.65
Frequency of occurrence of seizures at present	0.56	1.12	1.68	2.24	2.80	3.36	3.92	4.48	5.04
Cluster of seizures for several days, freedom from seizures for several weeks	0.77	1.54	2.31	3.08	3.85	NA	NA	NA	NA
Amount of EEG abnormality	1.57	3.14	4.71	6.28	7.85	NA	NA	NA	NA
Seizure patterns in EEG	0.09	0.18	0.27	0.36	0.45	NA	NA	NA	NA
	DISCRIMINATE FUNCTION II								
	1	2	3	4	5	6	7	8	9
Constant	-24.01								
Injuries during seizures	-0.02	-0.04	-0.06	-0.08	-0.10	NA	NA	NA	NA
Combination of seizures	0.06	0.12	0.18	0.24	0.30	0.36	0.42	0.48	0.54
Psychomotor seizures	6.25	12.50	NA	NA	NA	NA	NA	NA	NA
Duration of seizure disorder	2.92	5.84	8.76	11.68	14.60	17.52	20.44	23.36	26.28
Frequency of occurrence of seizures at present	1.02	2.04	3.06	4.08	5.10	6.12	7.14	8.16	9.18
Cluster of seizures for several days, freedom from seizures for several weeks	0.90	1.80	2.70	3.60	4.50	NA	NA	NA	NA
Amount of EEG abnormality	1.78	3.56	5.34	7.12	8.90	NA	NA	NA	NA
Seizure patterns in EEG	0.21	0.42	0.63	0.84	1.05	NA	NA	NA	NA

NA = Not Applicable

CONSTANTS AND WEIGHTS FOR PREDICTING OUTPATIENT TREATMENT RESULTS

	DISCRIMINATE FUNCTION I								
	1	2	3	4	5	6	7	8	9
Constant	-17.81								
Combination of seizures	-0.84	-1.68	-2.52	-3.36	-4.20	-5.04	-5.88	-6.72	-7.56
Clusters of seizures for several days, freedom from seizures for several weeks	-1.25	-2.50	-3.75	-5.00	-6.25	NA	NA	NA	NA
Injuries during seizures	1.25	2.50	3.75	5.00	6.25	NA	NA	NA	NA
Psychomotor seizures	9.48	18.96	NA	NA	NA	NA	NA	NA	NA
Duration of seizure disorder	1.92	3.84	5.76	7.68	9.60	11.52	13.44	15.36	17.28
Frequency of occurrence of seizures at present	1.35	2.70	4.05	5.40	6.75	8.10	9.45	10.80	12.15
Amount of EEG abnormality	2.46	4.92	7.38	9.84	12.30	NA	NA	NA	NA
Seizure patterns in EEG	0.40	0.80	1.20	1.60	2.00	NA	NA	NA	NA
	DISCRIMINATE FUNCTION II								
	1	2	3	4	5	6	7	8	9
Constant	-30.28								
Combination of seizures	-0.27	-0.54	-0.81	-1.08	-1.35	-1.62	-1.89	-2.16	-2.43
Clusters of seizures for several days, freedom from seizures for several weeks	-1.57	-3.14	-4.71	-6.28	-7.85	NA	NA	NA	NA
Injuries during seizures	1.92	3.84	5.76	7.68	9.60	NA	NA	NA	NA
Psychomotor seizures	11.37	22.74	NA	NA	NA	NA	NA	NA	NA
Duration of seizure disorder	2.60	5.20	7.80	10.40	13.00	15.60	18.20	20.80	23.40
Frequency of occurrence of seizures at present	1.66	3.32	4.98	6.64	8.30	9.96	11.62	13.28	14.94
Amount of EEG abnormality	2.20	4.40	6.60	8.80	11.00	NA	NA	NA	NA
Seizure patterns in EEG	1.36	2.72	4.08	5.44	6.80	NA	NA	NA	NA

NA = Not Applicable

DISTRIBUTION OF AGE AT DEATH FOR ALL INDIVIDUALS AND FOR
THOSE WITH EPILEPSY MENTIONED ON DEATH CERTIFICATES*

		<i>Total</i>	<i>Infants</i>	<i>1-4</i>	<i>5-9</i>	<i>10-14</i>	<i>15-19</i>	<i>20-24</i>	<i>25-29</i>	<i>30-34</i>	<i>35-39</i>
1960	Total Deaths	67,912	4,702	716	389	302	475	524	529	760	1,110
	Epilepsy Deaths	81	1	7	7	2	3	6	8	11	10
1961	Total Deaths	67,375	4,604	677	403	273	478	527	454	718	1,143
	Epilepsy Deaths	74	—	4	3	4	3	5	6	5	5
1962	Total Deaths	70,049	4,367	609	372	319	450	521	483	700	1,084
	Epilepsy Deaths	95	2	2	1	4	8	5	5	9	9
1963	Total Deaths	72,438	4,150	628	398	309	580	596	542	733	1,204
	Epilepsy Deaths	91	—	5	2	5	7	5	14	8	9
1964	Total Deaths	72,129	4,043	615	408	340	652	691	544	790	1,161
	Epilepsy Deaths	97	1	3	4	8	5	9	4	10	15
1965	Total Deaths	73,665	3,936	596	375	338	672	705	574	753	1,186
	Epilepsy Deaths	109	1	1	4	7	3	6	10	6	14
1966	Total Deaths	74,596	3,751	629	418	403	760	766	631	710	1,142
	Epilepsy Deaths	119	1	3	2	2	5	7	9	5	11

* Courtesy Michigan Department of Public Health, Vital Records Section

<i>40-44</i>	<i>45-49</i>	<i>50-54</i>	<i>55-59</i>	<i>60-64</i>	<i>65-69</i>	<i>70-74</i>	<i>75-79</i>	<i>80-84</i>	<i>85-89</i>	<i>90 and Over</i>	<i>Median Age At Death</i>
1,742	2,604	3,650	4,855	6,184	8,130	9,049	8,595	6,934	4,386	2,276	68.3
5	6	3	5	—	5	1	1	2	3	—	35.2
1,070	2,570	3,501	4,703	6,142	7,976	9,070	8,629	6,980	4,481	2,376	68.7
12	3	3	7	4	3	3	1	2	—	1	40.8
1,742	2,623	3,698	4,891	6,306	8,331	9,484	9,189	7,390	4,809	2,591	69.1
7	9	8	8	5	4	3	3	2	—	1	41.8
1,822	2,679	3,858	5,000	6,422	8,437	9,792	9,550	7,873	5,041	2,734	69.3
7	6	2	6	2	4	5	3	1	—	—	34.7
1,875	2,751	3,874	5,086	6,451	8,038	9,760	9,620	7,881	4,943	2,606	69.2
6	3	4	5	3	4	7	3	2	1	—	36.5
1,948	2,824	3,936	5,126	6,576	8,163	9,721	10,010	8,234	5,180	2,812	69.5
7	11	7	8	8	6	4	2	4	—	—	41.8
1,983	2,851	4,108	5,253	6,736	7,986	9,905	10,061	8,502	5,160	2,896	69.5
21	8	9	10	8	5	4	3	2	4	—	43.5

CONSTANTS AND WEIGHTS FOR PREDICTING INTELLECTUAL LOSS

	DISCRIMINATE FUNCTION I								
	1	2	3	4	5	6	7	8	9
Constant	-79.22								
Frequency of occurrence of seizures at present	-1.60	-3.20	-4.80	-6.40	-8.00	-9.60	-11.20	-12.80	-14.40
Combination of seizures	5.85	11.70	17.55	23.40	29.25	35.10	40.95	46.80	52.65
Clusters of seizures for several days, freedom from seizures for several weeks	3.01	6.02	9.03	12.04	15.05	NA	NA	NA	NA
EEG background amplitude	2.92	5.84	8.76	11.68	14.60	17.52	20.44	23.36	26.28
Organic pathology suspected from psychological tests	9.22	18.44	27.66	36.88	46.10	NA	NA	NA	NA
Social factors contributing to illness	4.01	8.02	12.03	16.04	20.05	NA	NA	NA	NA
Full Scale IQ	0.99	(ALWAYS MULTIPLY BY ACTUAL FULL SCALE FIGURE)							

	DISCRIMINATE FUNCTION II								
	1	2	3	4	5	6	7	8	9
Constant	-71.88								
Frequency of occurrence of seizures at present	-1.53	-3.06	-4.59	-6.12	-7.65	-9.18	-10.71	-12.24	-13.77
Combination of seizures	4.99	9.98	14.97	19.96	24.95	29.94	34.93	39.92	44.91
Clusters of seizures for several days, freedom from seizures for several weeks	2.19	4.38	6.57	8.76	10.95	NA	NA	NA	NA
EEG background amplitude	3.20	6.40	9.60	12.80	16.00	19.20	22.40	25.60	28.80
Organic pathology suspected from psychological tests	9.79	19.58	29.37	39.16	48.95	NA	NA	NA	NA
Social factors contributing to illness	3.42	6.84	10.26	13.68	17.10	NA	NA	NA	NA
Full Scale IQ	0.93	(ALWAYS MULTIPLY BY ACTUAL FULL SCALE FIGURE)							

NA = Not Applicable

NAME: _____

LAFAYETTE CLINIC INPATIENT EPILEPSY REVIEW—DR. ROBIN

PROJECT 116

Coding Sheets

CARD 6

DATE OF EXAMINATION

1-2 ☐ ☐ Month4-5 ☐ ☐ Day7-8 ☐ ☐ Year

10 REFERRING SOURCE

- ☐ 0 Not recorded
- ☐ 1 Private M.D.
- ☐ 2 Agency other than State Hospital
- ☐ 3 Court or Police
- ☐ 4 Caro State Hospital
- ☐ 5 Northville State Hospital
- ☐ 6 Pontiac State Hospital
- ☐ 7 Ypsilanti State Hospital
- ☐ 8 Traverse City/Kalamazoo State Hospital
- ☐ 9 Lapeer/Plymouth State Hospital

12 RACE

- ☐ 0 Not recorded
- ☐ 1 White
- ☐ 2 Negro
- ☐ 3 American Indian
- ☐ 4 Mexican
- ☐ 5 Oriental
- ☐ 6
- ☐ 7
- ☐ 8
- ☐ 9

*14 BIRTH ORDER (For 9 or more, Code as 9)

☐

16 MAIN REASON FOR ADMISSION

- ☐ 0 Not recorded
- ☐ 1 Seizure control
- ☐ 2 Behavior control
- ☐ 3 Teaching or research purposes

* Asterisks indicate variables that were used in data analysis.

PATIENT'S GENERAL PAST MEDICAL AND PSYCHIATRIC HISTORY
(Check the boxes which are applicable. Code "0" for items on which no information.)

- 18 ☐ First born twin or other multiple birth
 20 ☐ Second, third, or fourth multiple birth
 *22 ☐ Delayed maturation
 24 ☐ Breathholding spells
 *26 ☐ Febrile convulsions (isolated episodes not directly related to current seizure disorder)
 *28 ☐ Nonfebrile convulsions (isolated episodes not directly related to current seizure disorder)
 30 ☐ Atypical spells: _____
 *32 ☐ Bedwetting
 *34 ☐ Neurotic symptoms: _____
 *36 ☐ Psychotic symptoms: _____
 *38 ☐ Personality disorder: _____
 *40 ☐ Police record: _____

FAMILY HISTORY

(Check the boxes which are applicable. If no family history available, cross out entire section. Omit questionable data.)

STILLBIRTHS

- *42 ☐ Male
 *44 ☐ Female
 46 ☐ Paternal side of family
 *48 ☐ Maternal side of family
 *50 ☐ More than one in family
 *52 ☐ Sex unknown

State relationship to patient: _____

EARLY INFANTILE DEATHS

- *54 ☐ Male
 *56 ☐ Female
 *58 ☐ Paternal side of family
 *60 ☐ Maternal side of family
 *62 ☐ More than one in family
 *64 ☐ Sex unknown

State relationship to patient: _____

- 00-70 ☐☐☐☐☐ Lafayette Clinic inpatient number
 71-78 Leave blank
 79-80 ☐☐ Card number

CARD 7

FAMILY HISTORY (cont'd)

TWINS

- *2 ☐ Male
*4 ☐ Female
*0 ☐ Paternal side of family
*8 ☐ Maternal side of family
*10 ☐ More than one in family

State relationship to patient:

CONGENITAL MALFORMATIONS

- 12 ☐ Male
14 ☐ Female
16 ☐ Paternal side of family
18 ☐ Maternal side of family
20 ☐ More than one in family

State relationship to patient:

Type: _____

MENTAL RETARDATION

- *22 ☐ Male
*24 ☐ Female
*26 ☐ Paternal side of family
*28 ☐ Maternal side of family
30 ☐ More than one in family

State relationship to patient:

BREATHHOLDING SPELLS

- 32 ☐ Male
34 ☐ Female
36 ☐ Paternal side of family
38 ☐ Maternal side of family
40 ☐ More than one in family

State relationship to patient:

INFANTILE FEBRILE CONVULSIONS

- *42 ☐ Male
*44 ☐ Female
*46 ☐ Paternal side of family
*48 ☐ Maternal side of family
50 ☐ More than one in family

State relationship to patient:

EPILEPSY

- *52 ☐ Male
 *54 ☐ Female
 *56 ☐ Paternal side of family
 *58 ☐ Maternal side of family
 *60 ☐ More than one in family

State relationship to patient:

61-65 Leave blank

66-78 Duplicate these columns from original data on Card 6.

79-80 ☐ ☐ Card number

CARD 8

OTHER NEUROLOGICAL DISEASES

- *2 ☐ Male
 *4 ☐ Female
 *6 ☐ Paternal side of family
 *8 ☐ Maternal side of family
 *10 ☐ More than one in family

State relationship to patient:

Specify type

DIABETES

- *12 ☐ Male
 *14 ☐ Female
 *16 ☐ Paternal side of family
 *18 ☐ Maternal side of family
 *20 ☐ More than one in family

State relationship to patient:

PSYCHIATRIC DISORDERS

- *22 ☐ Male
 *24 ☐ Female
 *26 ☐ Paternal side of family
 *28 ☐ Maternal side of family
 *30 ☐ More than one in family

State relationship to patient:

Specify type

SCHOOL HISTORY

*32 AMOUNT OF SCHOOLING

(Code only when education completed)

- ☐ 0 Not recorded
- ☐ 1 No formal schooling
- ☐ 2 Attended grade school; did not complete 8th grade
- ☐ 3 Attended grade school; graduated 8th grade
- ☐ 4 Attended high school; not graduated
- ☐ 5 Graduated from high school
- ☐ 6 Attended college; not graduated
- ☐ 7 Graduated from college; include in "college" any recognized formal post-high school
- ☐ 8 Graduate school
- ☐ 9 Doctoral degree

*34 ATTENDING OR ATTENDED SPECIAL SCHOOL

- ☐ 0 Not recorded
- ☐ 1 No
- ☐ 2 Yes

*36 ATTENDING OR ATTENDED REGULAR SCHOOL

- ☐ 0 Not recorded
- ☐ 1 No
- ☐ 2 Yes

*38 AVERAGE GRADES

- ☐ 0 Not recorded
- ☐ 1 Unsatisfactory and grade failure
- ☐ 2 Unsatisfactory but no grade failure
- ☐ 3 Average
- ☐ 4 Somewhat above average
- ☐ 5 Superior or honor student

*40 USE OF ALCOHOL BY PATIENT

- ☐ 0 Not recorded
- ☐ 1 Abstinent
- ☐ 2 Social drinker
- ☐ 3 One or more drinks per day
- ☐ 4 Chronic alcoholism without DTs or Korsakoff's syndrome
- ☐ 5 Chronic alcoholism with DTs or Korsakoff's syndrome

TYPE (S) OF CLINICAL SEIZURES PRESENT

- *42 ☐ *Grand Mal Seizure, nonfocal:* No aura, loss of consciousness, falling, tonic phase, clonic phase, followed by sleep, some disorientation and memory loss.
- *44 ☐ *Grand Mal Seizure, focal onset:* Either subjective aura, or head and neck turning to one side, or twitching of one side of the face or part of an extremity, loss of consciousness, tonic phase, clonic phase, sleep; on recovery possible weakness of one extremity or pronounced disorientation and memory loss.

- *46 ☐ *Grand Mal Seizure* but history inadequate to differentiate between focal or nonfocal onset.
- *48 ☐ *Grand Mal Variant Seizure—Nonfocal*: Loss of consciousness, falling, some twitching or shaking, no well-defined tonic and clonic phase like in classical grand mal, there may or may not be sleep afterward.
- *50 ☐ *Grand Mal Variant—Focal onset*: Subjective aura, or head and neck turning to one side or twitching of one side of the face or twitching of one part of an extremity, loss of consciousness, falling, no tonic and clonic phase as in classical grand mal; some twitching or shaking, or merely generalized rigidity, there may or may not be sleep afterward.
- *51 ☐ *Minor Motor Focal Seizure*: Consciousness may be retained or lost, no falling, patient feels weak and sits down or remains standing, twitching of one or more extremities, or twitching of the face, no Jacksonian march, may or may not sleep afterward.
- *52 ☐ *Minor Motor Nonfocal Seizure*: Consciousness clouded, bilaterally symmetrical tonic extension or flexion of arms. There may be twitching but no definite jerking. May be confused afterwards, usually does not sleep. Retains posture or staggers.
- 53 ☐ *Minor Motor or Sensory Seizure with Jacksonian March*: Twitching and/or numbness starting in one discreet area (face, hand, leg) spreading in typical Jacksonian type of march.
- *54 ☐ *Automatism (Psychomotor)*: There may or may not be subjective aura, consciousness lost, posture retained, purposeless coordinated behavior which is inappropriate at that moment; when coming out of the seizure, patient is disoriented and may be belligerent, may or may not sleep, but sleep unlikely.
- 55 ☐ *Psychic or Sensory Seizure*: Subjective sensation on part of patient with no overt motor manifestations which occur suddenly and are not related to the situation the patient finds himself in; no loss of consciousness, no falling, no longer than about 15 minutes—usually only 1 to 2 minutes—usually no sleep afterward.
- 56 ☐ *Confusional State*: Confused and/or disoriented behavior lasting from several minutes to several hours, which is not preceded nor followed by a seizure.
- *57 ☐ *Absence*: Consciousness lost, staring expression, posture unchanged, continues with activities as soon as seizure is over at precisely the point at which he had been interrupted as a result of the seizure, and no sleep after seizure.
- *58 ☐ *Absence with features of slight automatism*: Some chewing or other slight repetitive activity.
- *59 ☐ *Absence with features of mild myoclonic activity*: Blinking of eyelids, slight jerking of head or hands.
- 60 ☐ *Myoclonic Seizure*: Clouding of consciousness, posture may be retained, twitching of eyelids, back and fro movements of the head (nodding type), and bilaterally symmetrical jerking of arms and/or legs, no after-effects.
- *61 ☐ *Myoclonic Jerks*: Isolated irregular jerks of one extremity with no change in consciousness and no loss of posture.

- *62 ☐ *Akinetic Seizure*: Loss of consciousness and postural tone, patient gets up immediately after falling and has no other symptoms.
- *63 ☐ *Temporal Lobe Components to Seizure Pattern*: (Check this box regardless how many other boxes have been checked if it is applicable—it serves as a summary column).
- 64 ☐ *Difficult to Classify*
- 65 Leave blank
- 66-78 Duplicate these columns from original data on Card 6.
- 79-80 ☐ ☐ ☐ Card number

CARD 9

ONSET OF SEIZURES

AGE AT TIME OF FIRST SEIZURE

- *1-3 ☐ ☐ ☐ Months

AGE AT ONSET OF RECURRENT SEIZURES

- *5-7 ☐ ☐ ☐ Months

TOTAL DURATION OF ILLNESS PRIOR TO HOSPITALIZATION

- *9-10 ☐ ☐ Years

REMISSION IN SEIZURE DISORDER

- *12-13 ☐ ☐ Years

*15 COMBINATION OF SEIZURES

(The even numbers are to be checked if the patient has focal grand mal seizures and sometimes in addition auras only. The aura would then not represent a second seizure type. The scale would be marked on number 2. If there are focal grand mal seizures and psychomotor automatisms and auras only, the scale would be marked at 4, etc.)

- ☐ Not recorded
- ☐ 1 seizure type only
- ☐ 2 seizure types
- ☐ 3 seizure types
- ☐ 4 seizure types
- ☐ More than four seizure types

ETIOLOGY OF SEIZURES

- *17 ☐ UNKNOWN (Check only if none of the other columns apply and adequate history available—place "0" in box if history inadequate.)

***19 PRENATAL OR PERINATAL INJURY**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Questionable
- ☐ 3 Probable
- ☐ 4 Very likely
- ☐ 5 Definite

***21 POSTNATAL HEAD INJURY**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 *Mild*: marked bruise or laceration, but no aftereffects.
- ☐ 3 *Moderate*: marked bruise or laceration with or without skull fracture, no unconsciousness, but may have been momentarily dazed. Some aftereffects like vomiting or dizziness for a few days.
- ☐ 4 *Marked*: injury associated with unconsciousness for a period of less than 5 minutes.
- ☐ 5 *Severe*: injury associated with unconsciousness for more than 5 minutes.

***23 CEREBRAL INFECTION**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Questionable
- ☐ 3 Probable
- ☐ 4 Very likely
- ☐ 5 Definite

***25 OTHER SIGNIFICANT EXTERNAL CAUSES**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Questionable
- ☐ 3 Probable
- ☐ 4 Very likely
- ☐ 5 Definite. Specify: _____

***27 MIXED HEREDITARY AND EXTERNAL CAUSE**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Questionable
- ☐ 3 Probable
- ☐ 4 Very likely
- ☐ 5 Definite

***20 HEREDITARY CAUSE ONLY**

(A family includes 2nd cousins, great-aunts, and great-uncles, but not great-grandparents. If column 27 is checked this column becomes "0"—not recorded.)

- ☐ 5 Not recorded
- ☐ 1 *Absent*: no family history of infantile convulsions or epilepsy.
- ☐ 2 *Questionable*: one other family member had isolated infantile convulsions.
- ☐ 3 *Probable*: several other family members had infantile convulsions or one close relative had epilepsy.
- ☐ 4 *Very likely*: several family members had infantile convulsions and one other had epilepsy.
- ☐ 5 *Definite*: two or more family members had epilepsy.

SPECIFIC SEIZURE HISTORY**31 MOST DIFFICULT PROBLEM**

- ☐ 0 Not recorded
- ☐ 1 Major seizures
- ☐ 2 Minor seizures

TYPE OF SEIZURE

- 33-34 ☐ ☐ (Indicate the type of seizure by noting the IBM column numbers which appear on Card 8, columns 42-62.)

CURRENT SEIZURE PRESENT SINCE FIRST YEAR OF LIFE

- *36-38 ☐ ☐ ☐ Months

***40 DURATION OF THIS PARTICULAR SEIZURE TYPE**

- ☐ 0 Not recorded
- ☐ 1 Less than 1 month
- ☐ 2 1-2 months
- ☐ 3 3-6 months
- ☐ 4 7-11 months
- ☐ 5 1-3 years
- ☐ 6 4-6 years
- ☐ 7 7-9 years
- ☐ 8 10-15 years
- ☐ 9 More than 15 years

***42 FREQUENCY OF MAXIMAL OCCURRENCE OF SEIZURES**

- ☐ 0 Not recorded
- ☐ 1 Less than once a year
- ☐ 2 About once a year
- ☐ 3 2-3 seizures a year
- ☐ 4 4-6 seizures a year
- ☐ 5 7-12 seizures a year
- ☐ 6 Once a month
- ☐ 7 2-3 a month
- ☐ 8 Once a week
- ☐ 9 Several a week

***44 FREQUENCY OF OCCURRENCE AT PRESENT**

- ☐ 0 Not recorded
- ☐ 1 Less than once a year
- ☐ 2 About once a year
- ☐ 3 2-3 seizures a year
- ☐ 4 4-6 seizures a year
- ☐ 5 7-12 seizures a year
- ☐ 6 Once a month
- ☐ 7 2-3 a month
- ☐ 8 Once a week
- ☐ 9 Several a week

***46 IN REMISSION AT PRESENT**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Remission lasted 1 year
- ☐ 3 Remission lasted 1-2 years
- ☐ 4 Remission 2-3 years
- ☐ 5 Remission 3-5 years
- ☐ 6 Remission 5-7 years
- ☐ 7 Remission 7-10 years
- ☐ 8 Remission 10-15 years
- ☐ 9 More than 15 years

***48 REMISSION IN THE PAST**

- ☐ 0 Not recorded
- ☐ 1 Never
- ☐ 2 Up to 1 year
- ☐ 3 1-2 years
- ☐ 4 3-5 years
- ☐ 5 6-8 years
- ☐ 6 9-11 years
- ☐ 7 12-14 years
- ☐ 8 15-20 years
- ☐ 9 More than 20 years

***50 CLUSTERS OF SEIZURES IN ONE DAY**

- ☐ 0 Not recorded
- ☐ 1 Never
- ☐ 2 Rarely
- ☐ 3 Occasionally
- ☐ 4 Frequently
- ☐ 5 Usually

***52 STATUS EPILEPTICUS**

- ☐ 0 Not recorded
- ☐ 1 Never
- ☐ 2 Rarely
- ☐ 3 Occasionally
- ☐ 4 Frequently
- ☐ 5 Usually

***54 CLUSTERS OF SEIZURES OVER SEVERAL DAYS THEN FREEDOM FOR WEEKS**

- ☐ 0 Not recorded
- ☐ 1 Never
- ☐ 2 Rarely
- ☐ 3 Occasionally
- ☐ 4 Frequently
- ☐ 5 Usually

***56 FREQUENCY OF OCCURRENCE OF AURA**

- ☐ 0 Not recorded
- ☐ 1 Never
- ☐ 2 Rarely
- ☐ 3 Occasionally
- ☐ 4 Frequently
- ☐ 5 Usually

57-65 Leave blank

66-78 Duplicate these columns from original data on Card 6.

79-80 ☐ 0 ☐ 9 Card number

CARD 10

TYPE OF AURA

- 2 ☐ Diffuse myoclonic jerks
- 4 ☐ Motor manifestations—right
- 6 ☐ Motor manifestations—left
- 8 ☐ Somatic sensory manifestations—right
- 10 ☐ Somatic sensory manifestations—left
- 12 ☐ Abdominal sensations
- 14 ☐ Vertigo
- 16 ☐ "Dizziness" other than vertigo
- 18 ☐ Déjà vue
- 20 ☐ Visual, auditory, or olfactory hallucinations
- 22 ☐ Fear
- 24 ☐ Difficulty speaking

***26 INJURIES SUSTAINED DURING ATTACK**

- ☐ 0 Not recorded
- ☐ 1 Never
- ☐ 2 Rarely
- ☐ 3 Occasionally
- ☐ 4 Frequently
- ☐ 5 Usually

***28-29 LENGTH OF HOSPITALIZATION**

☐ ☐ Weeks

***31 NUMBER OF ADMISSIONS**

☐

- *33-35 NUMBER OF MAJOR SEIZURES IN HOSPITAL WHILE ON MEDICATION

- *37-39 NUMBER OF MINOR SEIZURES WHILE ON MEDICATION

- 41 SEIZURES RELATED TO MENSES WHILE IN HOSPITAL

<input type="text"/>	Not recorded
<input type="text"/>	No
<input type="text"/>	Suggestive
<input type="text"/>	Definitely

- *43 SEIZURES RELATED TO LEAVE OF ABSENCE

<input type="text"/>	Not recorded
<input type="text"/>	No
<input type="text"/>	Suggestive
<input type="text"/>	Definitely

- *45 SEIZURES RELATED TO TIME OF DAY

<input type="text"/>	Not recorded
<input type="text"/>	No
<input type="text"/>	Suggestive
<input type="text"/>	Definitely

- 47-48 IF RELATED TO TIME OF DAY, MOST COMMON TIME OF OCCURRENCE (Based on 24 hours)

- *50 SEIZURES OCCURRED IN WAKING HOURS ONLY

<input type="text"/>	Not recorded
<input type="text"/>	No
<input type="text"/>	Yes

- *52 SEIZURES OCCURRED DURING SLEEP ONLY

<input type="text"/>	Not recorded
<input type="text"/>	No
<input type="text"/>	Yes

- *54 SEIZURES OCCURRED DURING WAKING AND SLEEPING

<input type="text"/>	Not recorded
<input type="text"/>	No
<input type="text"/>	Yes

LATERALITY OF CEREBRAL DISTURBANCE

- *56 RIGHT:

<input type="text"/>	Not recorded
<input type="text"/>	EEG only
<input type="text"/>	EEG and seizure pattern or neurological examination
<input type="text"/>	EEG, seizure pattern and neurological examination

*58 LEFT:

- ☐ 0 Not recorded
- ☐ 1 EEG only
- ☐ 2 EEG and seizure pattern or neurological examination
- ☐ 3 EEG, seizure patterns and neurological examination

*60 BILATERAL CEREBRAL INVOLVEMENT

(based on EEG, seizure patterns or neurological examination)

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Suggestively present
- ☐ 3 Definitely present

*62 NO CEREBRAL DISTURBANCE (e.g. normal or Dys. I EEG, normal neurological, no focal seizure pattern)

- ☐ 0 Not recorded
- ☐ 1 Cerebral disturbance definitely present
- ☐ 2 Cerebral disturbance suggestively present
- ☐ 3 Cerebral disturbance definitely absent

*64 PNEUMOENCEPHALOGRAM DIFFUSE ATROPHY

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Suggestive dilatation of ventricles
- ☐ 3 Definite ventricular dilatation

66-78 Duplicate these columns from original data on Card 6.

79-80 ☐ 0 Card number

CARD 11

PNEUMOENCEPHALOGRAM FOCAL ATROPHY

*2 RIGHT:

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Suggestively present
- ☐ 3 Definitely present

*4 LEFT:

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Suggestively present
- ☐ 3 Definitely present

*6 PNEUMOENCEPHALOGRAM CORTICAL ATROPHY

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Suggestively present
- ☐ 3 Definitely present

8 ARTERIOGRAM FOCAL ABNORMALITY OF POSSIBLE ETIOLOGICAL SIGNIFICANCE

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Suggestively present
☐ 3 Definitely present

10 LATERALIZATION ON ARTERIOGRAM

- ☐ 0 Not recorded
☐ 1 No lateralization
☐ 2 Right side
☐ 3 Left side

*12 OBJECTIVE FINDINGS OF CEREBRAL PATHOLOGY IN NEUROLOGICAL EXAMINATION (apart from organic mental syndrome)

- ☐ 0 Not recorded
☐ 1 None
☐ 2 Mild
☐ 3 Moderate
☐ 4 Marked

PSYCHOLOGICAL TEST RESULTS

*14-16 VERBAL IQ

*18-20 PERFORMANCE IQ

*21-23 FULL SCALE IQ

*25 IMMATURITY

- ☐ 0 Not recorded
☐ 1 None
☐ 2 Mild
☐ 3 Moderate
☐ 4 Marked

*27 NEUROTIC TENDENCIES

- ☐ 0 Not recorded
☐ 1 None
☐ 2 Mild
☐ 3 Moderate
☐ 4 Marked

*29 PSYCHOTIC TENDENCIES

- ☐ 0 Not recorded
☐ 1 None
☐ 2 Mild
☐ 3 Moderate
☐ 4 Marked

*31 PERSONALITY DISTURBANCES

- ☐ 0 Not recorded
☐ 1 None
☐ 2 Mild
☐ 3 Moderate
☐ 4 Marked

*33 ORGANIC PATHOLOGY SUSPECTED

- ☐ 0 Not recorded
☐ 1 None
☐ 2 Mild
☐ 3 Moderate
☐ 4 Marked

SEIZURE PHOTOGRAPH ANALYSIS
 INITIAL SYMPTOMS CONSIST OF:

- 35 ☐ Marked motor restlessness
 37 ☐ Requests to stop test
 39 ☐ Complains of dizziness
 *41 ☐ Sits up
 43 ☐ Isolated body or head jerks
 45 ☐ Twitching of eyelids
 47 ☐ Facial twitching unlocalized
 49 ☐ Facial twitching, right
 51 ☐ Facial twitching, left
 53 ☐ Rhythmic blinking of eyelids
 55 ☐ Rhythmic head nodding
 57 ☐ Right arm twitching
 59 ☐ Right leg twitching
 61 ☐ Left arm twitching
 63 ☐ Left leg twitching
 64-65 Leave blank
 66-78 Duplicate these columns from original data on Card 6.
 79-80 ☐ ☐ Card number

CARD 12 INITIAL SYMPTOMS CONSIST OF:

- 2 ☐ Verbalization
 4 ☐ Vocalization
 6 ☐ Looking around confused
 8 ☐ Facial expression, bewildered
 10 ☐ Facial expression, frightened
 12 ☐ Facial expression, empty
 14 ☐ Facial expression, angry

- 16 ☐ Salivation
 *18 ☐ Chewing or swallowing or snacking
 *20 ☐ Head turns to right
 *22 ☐ Head turns to left
 24 ☐ Head swings to opposite side after initial turning
 *26 ☐ Head is retracted backwards
 28 ☐ Head is flexed towards chest
 *30 ☐ Eyes deviated to right
 *32 ☐ Eyes deviated to left
 34 ☐ Eyes swing to opposite side after initial turning
 *36 ☐ Eyes deviated upwards
 38 ☐ Eyes converge
 40 ☐ Extended right arm raised laterally
 42 ☐ Extended right arm raised anteriorly
 44 ☐ Extended right arm raised posteriorly
 46 ☐ Extended left arm raised laterally
 48 ☐ Extended left arm raised anteriorly
 50 ☐ Extended left arm raised posteriorly
 *52 ☐ Right arm flexed in elbow raised laterally
 *54 ☐ Right arm flexed in elbow raised anteriorly
 56 ☐ Right arm flexed in elbow raised posteriorly
 *58 ☐ Left arm flexed in elbow raised laterally
 *60 ☐ Left arm flexed in elbow raised anteriorly
 62 ☐ Left arm flexed in elbow raised posteriorly
 *64 ☐ Right arm held before face as if looking at head
 66-78 Duplicate these columns from original data on Card 6.
 79-80 ☐ ☐ Card number

CARD 13 INITIAL SYMPTOMS CONSIST OF: (*cont'd*)

- *2 ☐ Left arm held before face as if looking at hand
 4 ☐ Right arm held above head as if touching back of head
 6 ☐ Left arm held above head as if touching back of head
 8 ☐ Right leg raised in extension
 10 ☐ Left leg raised in extension
 *12 ☐ Right leg raised while flexed in knee
 *14 ☐ Left leg raised while flexed in knee
 16 ☐ Right leg raised higher than left
 18 ☐ Left leg raised higher than right
 *20 ☐ Jackknifing
 22 ☐ Hyperextension of entire body
 24 ☐ Entire body rolls to the right
 26 ☐ Entire body rolls to the left
 *28 ☐ Legs diverge

TONIC PHASE

- 30 ☐ Tonic facial contraction, right
 32 ☐ Tonic facial contraction, left
 *34 ☐ Right arm extended in elbow and wrist
 *36 ☐ Left arm extended in elbow and wrist
 *38 ☐ Right arm extended in elbow, flexed in wrist
 *40 ☐ Left arm extended in elbow, flexed in wrist
 42 ☐ Right arm crosses over left
 44 ☐ Left arm crosses over right
 46 ☐ Arms diverge
 *48 ☐ Fingers partially flexed, right hand
 *50 ☐ Fingers partially flexed, left hand
 *52 ☐ Fist, right
 *54 ☐ Fist, left
 *56 ☐ Right arm flexed in elbow
 *58 ☐ Left arm flexed in elbow
 59-65 Leave blank
 66-78 Duplicate these columns from original data on Card 6.
 79-80 ☐ ☐ 3 Card number

CARD 14—TONIC PHASE (cont'd)

- *2 ☐ Right leg extended
 *4 ☐ Left leg extended
 6 ☐ Right leg flexed at knee
 8 ☐ Left leg flexed at knee
 10 ☐ Right leg crossed over left
 12 ☐ Left leg crossed over right
 14 ☐ Legs diverge
 *16 ☐ Spontaneous Babinski's sign, right
 *18 ☐ Spontaneous Babinski's sign, left

POSTICTAL STATE

- 20 ☐ Aphasia
 22 ☐ Confused as to person
 24 ☐ Confused as to place
 26 ☐ Confused as to time
 28 ☐ Random uncontrolled movements
 30 ☐ Yelling
 32 ☐ Aggressive behavior
 34 ☐ Chewing, swallowing or smacking of lips
 *36 ☐ Automatism
 38 ☐ Sleep

ICTAL AUTOMATISM

40 DURATION

- ☐ 0 Not recorded
☐ 1 Less than 1 minute
☐ 2 Less than 1 to 2 minutes
☐ 3 Less than 2 to 3 minutes
☐ 4 Less than 3 to 5 minutes
☐ 5 Over 5 minutes

*42 ☐ Slight automatic acts, patient remains seated or lying.

*44 ☐ Marked rhythmic repetitive movements, patient remains seated or lying.

*46 ☐ Leaves chair, wanders about.

47-65 Leave blank

66-78 Duplicate these columns from original data on Card 6.

79-80 ☐ 1 ☐ 4 Card number

MICHIGAN EPILEPSY CENTER AND LAFAYETTE CLINIC
ELECTROENCEPHALOGRAPHY

Coding Sheets
Department 3

<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>

Patient number
EEG number

Name _____

PROJECT NAME _____

PHYSICIAN'S NAME _____

CARD 1

2-3 Physician number

RECORDING DATE

4-5 Month

6-7 Day

8-9 Year

AGE

10-11 Months

*12-18 Years

*14 SEX

Male

Female

15 HANDEDNESS

Not recorded

Right

Left

16 PATIENT STATUS

Not recorded

Outpatient

Inpatient

Other—Describe:

17 PLACE OF RESIDENCE

Not recorded

Greater Detroit

Outside of Detroit

18 REFERRING PERSON OR AGENCY

- ☐ 3 Not recorded
- ☐ 1 Self
- ☐ 2 Spouse
- ☐ 3 Patient, sibling, or child
- ☐ 4 More distant relative or friend
- ☐ 5 Social agency
- ☐ 6 Physician or other hospital (except State Hospital transfer)
- ☐ 7 State Hospital transfer
- ☐ 8 Court or police
- ☐ 9 Other—Describe:

*19 LAST SEIZURE

- ☐ 0 Not recorded
- ☐ 1 Within one day
- ☐ 2 1-3 days
- ☐ 3 4-7 days
- ☐ 4 8-14 days
- ☐ 5 More than 14 days

*20 ON ANTICONVULSANT MEDICATION

- ☐ 0 Not recorded
- ☐ 1 No
- ☐ 3 Yes

21-23 FAMILY NUMBER

24-28 NUMBER OF PROBAND

29 ANCESTRY

- ☐ 0 Not recorded
- ☐ 1 Father
- ☐ 2 Mother
- ☐ 3 Paternal grandfather
- ☐ 4 Paternal grandmother
- ☐ 5 Maternal grandfather
- ☐ 6 Maternal grandmother
- ☐ 7 Paternal aunt or uncle
- ☐ 8 Maternal aunt or uncle
- ☐ 9 Cousin

30 SIBLING ORDER

0
1
2
3
4
5
6
7
8
9

31 SIBLING TOTAL

0
1
2
3
4
5
6
7
8
9

32 CHILD

0
1
2
3
4
5
6
7
9

33 DRUG TREATMENT OF PATIENT OR RECENT ELECTRIC
CONVULSIVE TREATMENT

0	Not recorded
1	None
2	Drugs which do not affect EEG
3	Drugs which could affect EEG, but patient received inadequate doses
4	Tranquilizers in adequate amounts
5	Barbiturates in adequate amounts
6	Energizers or stimulants in adequate amounts
7	Combination of drugs in adequate amounts
8	Drugs which could affect EEG, adequate dose, but stopped for 48 hours
9	Electric convulsive treatment within the last two months

***34 EEG DIAGNOSIS**

- ☐ 0 Not recorded
- ☐ 1 Normal
- ☐ 2 Borderline
- ☐ 3 Mild abnormality
- ☐ 4 Moderate abnormality
- ☐ 5 Marked abnormality

***35 CONVULSIVE DISORDER SUSPECTED**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Questionable
- ☐ 3 Probable
- ☐ 4 Very likely, but not diagnostic
- ☐ 5 Diagnostic

***36 FOCAL LESION SUSPECTED**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Questionable
- ☐ 3 Probable
- ☐ 4 Very likely but not diagnostic
- ☐ 5 Diagnostic

ALPHA ACTIVITY

- 37 ☐ Present with eyes open (Score when not due to drowsiness)

- *38** ☐ 0 Not recorded
- ☐ 1 Absent
 - ☐ 2 Slight (less than 25% of recording time with eyes closed)
 - ☐ 3 Moderate (25-50% of recording time with eyes closed)
 - ☐ 4 Marked (50-75% of recording time with eyes closed)
 - ☐ 5 Excellent (75-100% of recording time with eyes closed)

THETA ACTIVITY

- *39** ☐ Present with eyes open (Score when not due to drowsiness)

- *40** ☐ 0 Not recorded
- ☐ 1 Absent
 - ☐ 2 Slight (less than 25% of recording time)
 - ☐ 3 Moderate (25-50% of recording time)
 - ☐ 4 Marked (50-75% of recording time)
 - ☐ 5 Massive (75-100% of recording time)

FAST ACTIVITY

- *41** ☐ 0 Not recorded
- ☐ 1 Absent
 - ☐ 2 Slight (less than 25% of recording time)
 - ☐ 3 Moderate (25-50% of recording time)
 - ☐ 4 Marked (50-75% of recording time)
 - ☐ 5 Massive (75-100% of recording time)

- 42 ☐ 0 Not recorded
☐ 1 Anterior head regions only
☐ 2 Diffuse

*43 DIFFUSE DELTA ACTIVITY

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Slight (less than 25% of recording time)
☐ 3 Moderate (25-50% of recording time)
☐ 4 Marked (50-75% of recording time)
☐ 5 Massive (75-100% of recording time)

DISCRETE RHYTHMIC SLOW WAVE FOCUS

- 44 ☐ Right frontal
 45 ☐ Left frontal
 46 ☐ Right motor
 47 ☐ Left motor
 48 ☐ Right parietal
 49 ☐ Left parietal
 50 ☐ Right occipital
 51 ☐ Left occipital
 *52 ☐ Right anterior temporal
 *53 ☐ Left anterior temporal
 *54 ☐ Right midtemporal
 *55 ☐ Left midtemporal
 56 ☐ Right posterior temporal
 57 ☐ Left posterior temporal

58 ENTIRE RIGHT HEMISPHERE

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present

*59 ENTIRE LEFT HEMISPHERE

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present

TYPE OF FOCUS

- *60 ☐ Theta
 *61 ☐ Complex discharge
 62 ☐ Sharp wave
 63 ☐ Spike
 64-65 Leave blank
 66-70 ☐ ☐ ☐ ☐ ☐ Patient number
 71-72 ☐ ☐ Project number
 73-74 ☐ ☐ Deck number
 75-76 ☐ ☐ Evaluation number
 77-78 ☐ 3 Department number
 79-80 ☐ 1 Card number

CARD 2

MORE THAN ONE RHYTHMIC FOCUS

- 2 ☐ Shifting focus
3 ☐ Mirror focus
4 ☐ Independent foci

DISCRETE RANDOM SLOW WAVE FOCUS

- 5 ☐ Right frontal
6 ☐ Left frontal
7 ☐ Right motor
8 ☐ Left motor
9 ☐ Right parietal
10 ☐ Left parietal
11 ☐ Right occipital
12 ☐ Left occipital
13 ☐ Right anterior temporal
14 ☐ Left anterior temporal
15 ☐ Right midtemporal
16 ☐ Left midtemporal
17 ☐ Right posterior temporal
18 ☐ Left posterior temporal

19 ENTIRE RIGHT HEMISPHERE

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present

20 ENTIRE LEFT HEMISPHERE

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present

21 INDEPENDENT FOCI

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present

GENERALIZED PAROXYSMAL ACTIVITY

(Express severity in frequency occurrence of bursts)

*22 ABORTIVE PAROXYSMS

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Slight
☐ 3 Moderate
☐ 4 Marked

***23 DEFINITE PAROXYSMS BUT NOT SPIKE WAVE**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Slight
- ☐ 3 Moderate
- ☐ 4 Marked

***24 ABORTIVE ATYPICAL SPIKE WAVES**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Slight
- ☐ 3 Moderate
- ☐ 4 Marked

***25 ATYPICAL SPIKE WAVES**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Slight
- ☐ 3 Moderate
- ☐ 4 Marked

***26 ATYPICAL SPIKE WAVES WITH MULTIPLE SPIKES**

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Slight
- ☐ 3 Moderate
- ☐ 4 Marked

27 CLASSICAL SPIKE WAVES (3 c/s)

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Present

28 ATYPICAL SEIZURE PATTERNS

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Present

29 HYPERSARRHYTHMIA

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Present

BACKGROUND RHYTHMS VOLTAGE

*30 LEFT

- ☐ 0 Not recorded
- ☐ 1 0-10 μ v
- ☐ 2 10-20
- ☐ 3 20-30
- ☐ 4 30-40
- ☐ 5 40-50
- ☐ 6 50-60
- ☐ 7 60-70
- ☐ 8 70-80
- ☐ 9 Above 80

*31 RIGHT

- ☐ 0 Not recorded
- ☐ 1 0-10 μ v
- ☐ 2 10-20
- ☐ 3 20-30
- ☐ 4 30-40
- ☐ 5 40-50
- ☐ 6 50-60
- ☐ 7 60-70
- ☐ 8 70-80
- ☐ 9 Above 80

32 SYMMETRY

- ☐ 0 Not recorded
- ☐ 1 More than 60 μ v lower on right
- ☐ 2 30-60 μ v lower on right
- ☐ 3 10-30 μ v lower on right
- ☐ 4 Up to 10 μ v lower on right
- ☐ 5 Background symmetrical
- ☐ 6 Up to 10 μ v higher on right
- ☐ 7 10-30 μ v higher on right
- ☐ 8 30-60 μ v higher on right
- ☐ 9 More than 60 μ v higher on right

MAIN BACKGROUND FREQUENCY

*33-34 ☐ ☐

HYPERVENTILATION

*35 DIFFUSE NONPAROXYSMAL BUILDUP

- ☐ 0 Not recorded
- ☐ 1 Absent
- ☐ 2 Slight
- ☐ 3 Moderate
- ☐ 4 Marked

36 ☐ Buildup persists for more than 1 minute

***37 DIFFUSE PAROXYSMAL BUILDUP**

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Slight
☐ 3 Moderate
☐ 4 Marked

38 ☐ Buildup persists for more than 1 minute

***39 FOCAL BUILDUP**

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present in resting record but increased with hyperventilation
☐ 3 Not present, brought out by hyperventilation

AREA OF FOCUS

- 40 ☐ Right frontal
41 ☐ Left frontal
42 ☐ Right motor
43 ☐ Left motor
44 ☐ Right parietal
45 ☐ Left parietal
46 ☐ Right occipital
47 ☐ Left occipital
48 ☐ Right anterior temporal
49 ☐ Left anterior temporal
50 ☐ Right midtemporal
51 ☐ Left midtemporal
52 ☐ Right posterior temporal
53 ☐ Left posterior temporal

54 SPECIFIC EPILEPTIC CHANGES

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present in resting record but increased with hyperventilation
☐ 3 Not present, brought out by hyperventilation

PHOTIC DRIVING RESPONSE**LOW FLASH RATES 1-7 f/s**

- *55** ☐ 0 Not recorded
☐ 1 None
☐ 2 Slight
☐ 3 Moderate
☐ 4 Marked
- 56 ☐ 0 Not recorded
☐ 1 Mostly subharmonic
☐ 2 Mostly fundamental
☐ 3 Mostly harmonic

- 57 ☐ 0 Not recorded
☐ 1 Occipital only
☐ 2 Parieto-occipital
☐ 3 Parieto-occipital-temporal
☐ 4 Motor-parieto-occipital-temporal
☐ 5 Diffuse

ALPHA RANGE 8-12 f/s

- *58 ☐ 0 Not recorded
☐ 1 None
☐ 2 Slight
☐ 3 Moderate
☐ 4 Marked
- 59 ☐ 0 Not recorded
☐ 1 Mostly subharmonic
☐ 2 Mostly fundamental
☐ 3 Mostly harmonic

- 60 ☐ 0 Not recorded
☐ 1 Occipital only
☐ 2 Parieto-occipital
☐ 3 Parieto-occipital-temporal
☐ 4 Motor-parieto-occipital-temporal
☐ 5 Diffuse

HIGH FREQUENCIES 13 f/s AND ABOVE

- *61 ☐ 0 Not recorded
☐ 1 None
☐ 2 Slight
☐ 3 Moderate
☐ 4 Marked
- 62 ☐ 0 Not recorded
☐ 1 Mostly subharmonic
☐ 2 Mostly fundamental
☐ 3 Mostly harmonic

- 63 ☐ 0 Not recorded
☐ 1 Occipital only
☐ 2 Parieto-occipital
☐ 3 Parieto-occipital-temporal
☐ 4 Motor-parieto-occipital-temporal
☐ 5 Diffuse

64-65 Leave blank

66-78 Duplicate these columns from original data on Card 1, Department 3.

79-80 ☐ 1 ☐ 2 Card number

CARD 8

OTHER PHOTIC RESPONSES

- 2 ☐ Photomyoclonic response
3 ☐ Excessive, spikey appearing driving response
4 ☐ Abortive paroxysms
5 ☐ Spike wave paroxysms without clinical accompaniments
6 ☐ Spike wave paroxysms with clinical accompaniments
7 ☐ Grand mal type buildup

MISCELLANEOUS PHENOMENA

- 8 ☐ Anterior displacement of alpha
9 ☐ Posterior head regions slowing
10 ☐ Alpha notching
11 ☐ Lambda waves
12 ☐ Spikey appearing record
13 ☐ Other—Describe _____

SLEEP

- *14 ☐ 0 Not recorded
☐ 1 Normal
☐ 2 Borderline
☐ 3 Mildly abnormal
☐ 4 Moderately abnormal
☐ 5 Markedly abnormal
- 15 ☐ 0 Not recorded
☐ 1 Induced
☐ 2 Spontaneous
- *16 SLEEP RECORD SHOWS MORE ABNORMALITY THAN WAKING RECORD
☐ 0 Not recorded
☐ 1 No
☐ 2 Yes
- *17 SLEEP RECORD SHOWS LESS ABNORMALITY THAN WAKING RECORD
☐ 0 Not recorded
☐ 1 No
☐ 2 Yes
- 18 STAGES RECORDED
☐ 0 Not recorded
☐ 1 Drowsiness only
☐ 2 V wave
☐ 3 Spindles
☐ 4 K complex spindles
☐ 5 Delta

MAIN TYPE OF ABNORMALITY

- 19 ☐ Spikey V wave or K complex
 20 ☐ Spindle asymmetry
 21 ☐ 14 and 6 c/s positive spikes
 *22 ☐ Focal sharp waves or spikes
 23 ☐ Focal slow waves
 24 ☐ Focal flattening
 *25 ☐ Paroxysmal abnormalities
 26 ☐ Atypical spike waves
 27 ☐ Classical spike waves
 28 ☐ Other—Describe: _____

AREA OF FOCAL ABNORMALITY

- 29 ☐ Right frontal
 30 ☐ Left frontal
 31 ☐ Right motor
 32 ☐ Left motor
 33 ☐ Right parietal
 34 ☐ Left parietal
 35 ☐ Right occipital
 36 ☐ Left occipital
 37 ☐ Right anterior temporal
 38 ☐ Left anterior temporal
 39 ☐ Right midtemporal
 40 ☐ Left midtemporal
 41 ☐ Right posterior temporal
 42 ☐ Left posterior temporal

MEGIMIDE STUDY

QUANTITY GIVEN (cc)

*43-44 ☐☐

*45 GENERALIZED SLOWING

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Slight
☐ 3 Moderate
☐ 4 Marked

*46 PAROXYSMAL RESPONSE

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Slight
☐ 3 Moderate
☐ 4 Marked

*47 FOCAL RESPONSE

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present in resting record but increased with drug
☐ 3 Not present, brought out by drug

AREA OF FOCAL RESPONSE

- 48 ☐ Right frontal
49 ☐ Left frontal
50 ☐ Right motor
51 ☐ Left motor
52 ☐ Right parietal
53 ☐ Left parietal
54 ☐ Right occipital
55 ☐ Left occipital
*56 ☐ Right anterior temporal
*57 ☐ Left anterior temporal
*58 ☐ Right midtemporal
*59 ☐ Left midtemporal
60 ☐ Right posterior temporal
61 ☐ Left posterior temporal

62 SEIZURE INDUCED

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 EEG seizure only
☐ 3 EEG and clinical seizure

63 ATTACK INDUCED WITHOUT EEG CHANGE

- ☐ 0 Not recorded
☐ 1 Absent
☐ 2 Present

64-65 Leave blank

66-78 Duplicate these columns from original data on Card 1, Department 3.

79-80 ☐ 3 Card number

MICHIGAN EPILEPSY CENTER AND LAFAYETTE CLINIC
EEG SEIZURE PATTERNS

Department 3

Coding Sheets

☐☐☐☐☐ Patient number NAME _____
PROJECT NAME _____
PHYSICIAN NAME _____

CARD 4

CLASSICAL 3 c/s SPIKE WAVE

2 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

3 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

CLASSICAL 3 c/s SPIKE WAVE WITH MULTIPLE SPIKES

4 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

5 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

CLASSICAL 3 c/s SPIKE WAVE WITH FOCAL ONSET

6 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

7 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

AREAS INVOLVED

- 8 ☐ Left fronto-motor
- 9 ☐ Right fronto-motor
- 10 ☐ Left motor-parietal
- 11 ☐ Right motor-parietal
- 12 ☐ Left parieto-occipital
- 13 ☐ Right parieto-occipital
- 14 ☐ Left anterior-midtemporal
- 15 ☐ Right anterior-midtemporal
- 16 ☐ Left mid-posterior temporal
- 17 ☐ Right mid-posterior temporal

4 c/s SPIKE WAVE

18 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

19 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

4 c/s SPIKE WAVE WITH MULTIPLE SPIKES

20 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

21 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

4 c/s SPIKE WAVE WITH FOCAL ONSET

22 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

23 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

AREAS INVOLVED

- 24 ☐ Left fronto-motor
- 25 ☐ Right fronto-motor
- 26 ☐ Left motor-parietal
- 27 ☐ Right motor-parietal
- 28 ☐ Left parieto-occipital
- 29 ☐ Right parieto-occipital
- 30 ☐ Left anterior-midtemporal
- 31 ☐ Right anterior-midtemporal
- 32 ☐ Left mid-posterior temporal
- 33 ☐ Right mid-posterior temporal

1-2 1/2 c/s SPIKE WAVE

34 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

35 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

1-2 1/2 c/s SPIKE WAVE WITH MULTIPLE SPIKES

36 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

37 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

1-2 1/2 c/s SPIKE WAVE WITH FOCAL ONSET

38 NUMBER OCCURRING IN RESTING RECORD

- ☐ 0 Not applicable
- ☐ 1 One
- ☐ 2 Two
- ☐ 3 Three
- ☐ 4 Four
- ☐ 5 Five
- ☐ 6 Six
- ☐ 7 Seven
- ☐ 8 Eight
- ☐ 9 Nine or more

39 AVERAGE DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 5 seconds
- ☐ 2 5-10 seconds
- ☐ 3 10-15 seconds
- ☐ 4 15-20 seconds
- ☐ 5 20-25 seconds
- ☐ 6 25-30 seconds
- ☐ 7 30-35 seconds
- ☐ 8 35-40 seconds
- ☐ 9 More than 40 seconds

AREAS INVOLVED

- 40 ☐ Left fronto-motor
- 41 ☐ Right fronto-motor
- 42 ☐ Left motor-parietal
- 43 ☐ Right motor-parietal
- 44 ☐ Left parieto-occipital
- 45 ☐ Right parieto-occipital
- 46 ☐ Left anterior-midtemporal
- 47 ☐ Right anterior-midtemporal
- 48 ☐ Left midposterior temporal
- 49 ☐ Right midposterior temporal

FOCAL TEMPORAL SEIZURE (a) STARTS WITH GRADUAL BUILDUP
OF DISCHARGES

AREA OF ONSET

- 50 ☐ Left anterior temporal
- 51 ☐ Left midtemporal
- 52 ☐ Left posterior temporal
- 53 ☐ Right anterior temporal
- 54 ☐ Right midtemporal
- 55 ☐ Right posterior temporal
- 56 ☐ Unclear

57 DURATION

- ☐ 0 Not applicable
- ☐ 1 Less than 30 seconds
- ☐ 2 31-60 seconds
- ☐ 3 1-3 minutes
- ☐ 3 3-6 minutes
- ☐ 6 More than 6 minutes

FOCAL TEMPORAL SEIZURE (b) STARTS WITH DIFFUSE BURST

AREA MOST INVOLVED SUBSEQUENTLY

- 58 ☐ Left anterior temporal
 59 ☐ Left midtemporal
 60 ☐ Left posterior temporal
 61 ☐ Right anterior temporal
 62 ☐ Right midtemporal
 63 ☐ Right posterior temporal
 64 ☐ Unclear

65 DURATION

- ☐ 0 Not applicable
☐ 1 Less than 30 seconds
☐ 3 31-60 seconds
☐ 3 1-3 minutes
☐ 4 3-6 minutes
☐ 5 More than 6 minutes

- 66-70 ☐☐☐☐ Patient number
 71-72 ☐☐ Project number
 73-74 ☐☐ Deck number
 75-76 ☐☐ Evaluation number
 77-78 ☐3 Department number
 79-80 ☐4 Card number

CARD 5

FOCAL TEMPORAL SEIZURE (c) STARTS WITH SUPPRESSION
OF ELECTRICAL ACTIVITY

AREA MOST INVOLVED SUBSEQUENTLY

- 2 ☐ Left anterior temporal
 3 ☐ Left mid-temporal
 4 ☐ Left posterior temporal
 5 ☐ Right anterior temporal
 6 ☐ Right mid-temporal
 7 ☐ Right posterior temporal
 8 ☐ Unclear

9 DURATION

- ☐ 0 Not applicable
☐ 1 Less than 30 seconds
☐ 3 31-60 seconds
☐ 3 1-3 minutes
☐ 4 3-6 minutes
☐ 3 More than 6 minutes

FOCAL SEIZURE OTHER THAN TEMPORAL, STARTS WITH GRADUAL
BUILDUP OF DISCHARGES

AREA OF ONSET

- 10 ☐ Left fronto-motor
11 ☐ Left motor-parietal
12 ☐ Left parieto-occipital
13 ☐ Right fronto-motor
14 ☐ Right motor-parietal
15 ☐ Right parieto-occipital

16 DURATION

- ☐ 0 Not applicable
☐ 1 Less than 30 seconds
☐ 2 31-60 seconds
☐ 3 1-3 minutes
☐ 4 3-9 minutes
☐ 5 More than 9 minutes

FOCAL SEIZURE OTHER THAN TEMPORAL, STARTS
WITH DIFFUSE BURST

AREA MOST INVOLVED SUBSEQUENTLY

- 17 ☐ Left fronto-motor
18 ☐ Left motor-parietal
19 ☐ Left parieto-occipital
20 ☐ Right fronto-motor
21 ☐ Right motor-parietal
22 ☐ Right parieto-occipital
23 ☐ Unclear

24 DURATION

- ☐ 0 Not applicable
☐ 1 Less than 30 seconds
☐ 2 31-60 seconds
☐ 3 1-3 minutes
☐ 4 3-9 minutes
☐ 5 More than 9 minutes

GRAND MAL SEIZURE, STARTS WITH GRADUAL BUILDUP
OF DISCHARGES

AREA OF ONSET

- 25 ☐ Left fronto-motor
 26 ☐ Right fronto-motor
 27 ☐ Left motor-parietal
 28 ☐ Right motor-parietal
 29 ☐ Left parieto-occipital
 30 ☐ Right parieto-occipital
 31 ☐ Left anterior-midtemporal
 32 ☐ Right anterior-midtemporal
 33 ☐ Left mid-posterior temporal
 34 ☐ Right mid-posterior temporal
 *35 ☐ Unclear
- *36 ☐ GRAND MAL SEIZURE, STARTS WITH DIFFUSE BURST

SEIZURE DIFFICULT TO CLASSIFY
ELECTROENCEPHALOGRAPHICALLY

*37 DURATION

- ☐ 0 Not applicable
☐ 1 Less than 10 seconds
☐ 2 11-30 seconds
☐ 3 31 seconds to 1 minute
☐ 4 1 minute to 3 minutes
☐ 5 More than 3 minutes

POSTICTAL SLOW WAVE FOCUS

- 38 ☐ Left fronto-motor
 39 ☐ Right fronto-motor
 40 ☐ Left motor-parietal
 41 ☐ Right motor-parietal
 42 ☐ Left parieto-occipital
 43 ☐ Right parieto-occipital
 44 ☐ Left anterior-midtemporal
 *45 ☐ Right anterior-midtemporal
 46 ☐ Left mid-posterior temporal
 *47 ☐ Right mid-posterior temporal
 *48 ☐ Generalized slowing

CLINICAL TYPE OF RECORDED SEIZURE

- *49 ☐ Grand mal focal
- *50 ☐ Grand mal nonfocal
- *51 ☐ Major motor not typical grand mal focal
- 52 ☐ Major motor not typical grand mal nonfocal
- 53 ☐ Minor motor focal
- 54 ☐ Minor motor nonfocal
- 55 ☐ Absence with minimal myoclonic element
- 56 ☐ Absence with marked myoclonic element
- 57 ☐ Myoclonic jerks
- *58 ☐ Automatism
- 59 ☐ Confusional state
- 60 ☐ Psychic
- 61 ☐ Sensory
- 62 ☐ Akinetic
- 63 ☐ Syncope
- 64 ☐ Psychogenic
- 65 ☐ Difficult to classify
- 66-78 Duplicate from Card 4, Department 3.
- 79-80 ☐ ☐ 5 Card number

REFERENCES

- AASS, F.; KAADA, B. R., and TORP, K. H.: The diagnostic and prognostic value of the initial electroencephalogram in children with convulsions. *Acta Paediat. (Stockholm)*, 45:335, (July) 1956.
- ABBOTT, JOHN A., and SCHWAB, ROBERT S.: Some clinical aspects of the normal electroencephalogram in epilepsy. *New Eng. J. Med.*, 238:457, 1948.
- ADIE, W. J.: Pyknolepsy: a form of epilepsy occurring in children with a good prognosis. *Brain*, 47:96, 1924.
- AJMONE-MARSAN, COSIMO, and BALDWIN, MAITLAND: Electrocorcicography. In *Temporal Lobe Epilepsy*, Baldwin, M., and Bailey, P. (Eds.). Springfield, Illinois, Thomas, 1958.
- ALAJOUANINE, TH.; MAISONNET, J., and PETIT-DUTAILLIS, D.: Suites éloignées de la trépanation du crâne pour lésions traumatiques. *J. Chir. (Paris)*, 32:397, 1928.
- ALSTROEM, CARL HENRY: A study of epilepsy in its clinical, social and genetic aspects. *Munksgaard* (Copenhagen), 1950.
- ARIEFF, ALEX J., and YACORZYNSKI, G. K.: Deterioration of patients with organic epilepsy. *J. Nerv. Ment. Dis.*, 96:49, 1942.
- ARIEFF, ALEX J.: Twelve-year resumé in a clinic for epilepsy. *Dis. Nerv. Syst.*, 12:19, 1951.
- ARNOLD, O. H.: Epilepsie. Eine statistische Studie am Material einer Epileptikerambulanz. *Wien. Z. Nervenheilk.*, 9:359, 1954.
- ASCROFT, P. B.: Traumatic epilepsy after gunshot wounds of the head. *Brit. Med. J.*, 1:739, (May) 1941.
- BAILEY, PERCIVAL, and GIBBS, FREDERIC A.: The surgical treatment of psychomotor epilepsy. *JAMA*, 145:365, (Feb) 1951.
- BAILEY, PERCIVAL; GREEN, JOHN R.; AMADOR, LUIS, and GIBBS, FREDERIC A.: Treatment of psychomotor states by anterior temporal lobectomy: a report of progress. *Proc. Ass. Res. Nerv. Ment. Dis.*, 31:341, 1953.
- BAIRD, H. W., III, and BONOFKY, L. G.: Infantile myoclonic seizures. *J. Pediat.*, 50:332, 1957.
- BALDWIN, MAITLAND, and BAILEY, PEARCE (Eds.): *Temporal Lobe Epilepsy*. Springfield, Illinois, Thomas, 1958.
- BAMBERGER, P., and MATTHES, A.: *Anfalle im Kindesalter*. Basel (Schweiz), Karger, and White Plains, N.Y., Pliebig, 1959.
- BARNES, MARGARET R., and FETTERMAN, JOSEPH L.: Mentality of dispensary epileptic patients. *Arch. Neurol. Psychiat.*, 40:903, 1938.
- BAUMM, HANS: Erfahrungen ueber Epilepsie bei Hirnverletzten. *Z. Ges. Neurol. Psychiat.*, 127:279, 1930.
- BERGAMINI, L.; BRAM, S.; BROGLIA, S., and RICCIO, A.: L'insorgenza tardiva di crisi grande male nel piccolo male puro: Studio catamnestico di 78 casi. *Schweiz. Arch. Neurol. Psychiat.*, 96:306, 1965.

- BERGEMANN, EVA: Ueber das spaetere Schicksal von Kindern mit Krampfaellen. *Msehr. Kinderheilk.*, 65:116, 1936.
- BLOOM, DAVID; JASPER, HERBERT, and RASMUSSEN, THEODORE: Surgical therapy in patients with temporal lobe seizures and bilateral EEG abnormality. *Epilepsia*, 1:351, 1960.
- BOWER, B. D., and JEA VONS, P. M.: Infantile spasms and hypsarrhythmia. *Lancet*, 1:605, 1959.
- BOWER, B. D., and JEA VONS, P. M.: The effect of Corticotrophin and Prednisolone on infantile spasms with mental retardation. *Arch. Dis. Child.*, 36:23, 1961.
- BRAIN, W. RUSSELL: *Diseases of the Nervous System*, 4th ed. London, Geoffrey Cumberledge, Oxford U.P., 1951.
- BRAY, P. F.: The influence of adrenal steroids and corticotrophin on massive myoclonic seizures of infancy. *J. Pediat.*, 32:169, (Aug.) 1963.
- BREG, W. ROY, and YANNET, HERMAN: The child in a convulsion. *Pediat. Clin. N. Amer.*, 9:101, 1962.
- BRIDGE, EDWARD M.: *Epilepsy and Convulsive Disorders in Children*, 1st ed. New York, McGraw, 1949.
- BRUN, R.: Zur Frage der posttraumatischen Epilepsien Vorbelastung und Katamnesen. *Zbl. Ges. Neurol. Psychiat.*, 76:360, 1955.
- BURKE, JOYCE B.: The prognostic significance of neonatal convulsions. *Arch. Dis. Child.*, 29:342, 1954.
- CAVENESS, WILLIAM F., and LISS, HENRY R.: Incidence of post-traumatic epilepsy. *Epilepsia*, 2:123, 1961.
- CAVENESS, WILLIAM F.: Onset and cessation of fits following craniocerebral trauma. *J. Neurosurg.*, 20:570, 1963.
- CHAO, DORA HSI-CHIH; TAYLOR, FRED M., and DRUCKMAN, RALPH: Massive spasms. *J. Pediat.*, 50:670, 1957.
- CHAO, DORA HSI-CHIH; DRUCKMAN, RALPH, and KELLAWAY, PETER: *Convulsive Disorders of Children*. Philadelphia, Saunders, 1958.
- CHAO, DORA HSI-CHIH: Seizures in infancy and early childhood. *Med. Clin. N. Amer.*, 42:399, 1958.
- CHAUDHRY, M. R., and POND, D. A.: Mental deterioration in epileptic children. *J. Neurol. Psychiat.*, 24:213, 1961.
- CLEMMESEN, CARL, and MOLTKE, MARIE LOUISE: Die Prognose der Kinder-epilepsie. Abstracted in *Zbl. Ges. Neurol. Psychiat.*, 75:425, 1935.
- CLUSZ, K.: Ueber Dauererfolge der operativen Behandlung der traumatischen Jackson'schen Epilepsie. *Beitr. Klin. Chir.*, 66:225, 1910.
- COBB, WILLIAM, and PAMPIGLIONE, GIUSEPPE: The electroencephalogram of hemiplegic patients treated by hemispherectomy. Abstracted in *Electroenceph. Clin. Neurophysiol., Suppl.* 3:62, 1953.
- COLLINS, A. LOUISE, and LENNOX, WILLIAM G.: The intelligence of 300 private epileptic patients. *Proc. Ass. Res. Nerv. Ment. Dis.*, 26:586, 1947.

- COLLINS, A. LOUISE: Epileptic intelligence. *J. Consult. Psychol.*, 15:392, 1951.
- COOPER, J. E.: Epilepsy in a longitudinal survey of 5,000 children. *Brit. Med. J.*, 1:1020, 1965.
- COURJON, JEAN, and COTTE, MME.: L'épilepsie avant trois ans. *Rev. Neurol. (Paris)*, 99:68, 1958.
- CRAIG, W. S., and MACKINNON, JOAN M.: Convulsions in infancy and childhood—Part II. *Brit. Med. J.*, No. 5433:499, 1965.
- CREDNER, LENE: Klinische und soziale Auswirkungen von Hirnschädigungen. *Z. Ges. Neurol. Psychiat.*, 126:721, 1930.
- CURRIER, ROBERT D.; KOOL, KENNETH A., and SAIDMAN, LAWRENCE, J.: Prognosis of "pure" petit mal: A follow-up study. *Neurology*, 13:959, (Nov.) 1963.
- DANIELSEN, J.: Infantile spasms and hypsarhythmia treated with corticosteroids. *Acta Neurol. Scand.*, 41:489, 1965.
- DAWSON, SHEPHERD, and CONN, J. C. M.: The intelligence of epileptic children. *Arch. Dis. Child.*, 4:142, 1929.
- DEKABAN, ANATOLE: Idiopathic epilepsy in early infancy. *Amer. J. Dis. Child.*, 100:181, (Aug.) 1960.
- DELAVELEYE, R., and SAUVEUR, J.-P.: Relation entre divers aspects de l'épilepsie infantile et le niveau intellectuel. *Rev. Neuropsychiat.*, 12:217, 1964.
- DENNERLL, R. D.; DEN BROEDER, JOAN, and SOKOLOV, S. L.: WISC and WAIS factors in children and adults with epilepsy. *J. Clin. Psychol.*, 20:236, (Apr.) 1964.
- DENNERLL, R. D.: Cognitive deficits and lateral brain dysfunction in temporal lobe epilepsy. *Epilepsia*, 5:177, 1964.
- DENNY-BROWN, D.: The sequelae of war head injuries. *New Eng. J. Med.*, 227:771, (Nov.) 1942.
- DENNY-BROWN, D.: The clinical aspects of traumatic epilepsy. *Amer. J. Psychiat.*, 100:585, (Mar.) 1944.
- DIXON, W. J. (Ed.): BMD-05M, Bio-Medical Computer Programs, Health Sciences Computing Facility, U.C.L.A., (Sept.) 1965.
- DOOSE, H., and SCHEFFNER, D.: Zur Therapie und Prognose der Absencen. *Med. Welt*, 1763, 1962.
- DOOSE, H.: *Verlaufsbilder der Petit Mal-Epilepsie im Kindesalter*. Kiel, Habilitationsschrift, 1963.
- DOOSE, H.; PETERSEN, C. E.; VOELZKE, E., and HERZBERGER, E.: Fieberkrämpfe und Epilepsie, I. Aetiologie, klinisches Bild und Verlauf der sogenannten Infekt- oder Fieberkrämpfe, II. Elektrencephalographische Verlaufsuntersuchungen bei sogenannten Fieber- oder Infektkrämpfen. *Arch. Psychiat. Nervenkr.*, 208:400, (July/August) 1966.
- DUMERMUTH, G.: Ueber das Syndrom der Blitz-Nick-Salaam-Krämpfe und

- seine Behandlung mit ACTH und Hydrocortison. *Helv. Paediat. Acta*, 16:244, 1961.
- EKHOLM, ERIK, and NIEMINEVA, KALEVI: On convulsions in early childhood and their prognosis: An investigation with follow-up examinations of patients treated for convulsions at the Children's Clinic of Helsinki University. *Acta Paediat.*, 39:481, 1950.
- EVANS, JOHN H.: Post-traumatic epilepsy. *Neurology*, 12:665, 1962.
- EVANS, JOHN H.: The significance of early post-traumatic epilepsy. *Neurology*, 13:207, 1963.
- FABING, HOWARD D., and BARROW, ROSCOE L.: Medical progress in treating epilepsy and the need for reform of laws affecting epileptics. *Epilepsia*, 3:92, 1954.
- FALCONER, MURRAY A.; HILL, DENIS; MEYER, ALFRED, and WILSON, JOHN L.: Clinical, radiological, and EEG correlations with pathological changes in temporal lobe epilepsy and their significance in surgical treatment. In *Temporal Lobe Epilepsy*, Baldwin, M., and Bailey, P. (Eds.). Springfield, Illinois, Thomas, 1958.
- FALCONER, M. A., and SERAFETINIDES, E. A.: A follow-up study of surgery in temporal lobe epilepsy. *J. Neurol., Neurosurg. Psychiat.*, 26:154, 1963.
- FALCONER, M. A.: Die chirurgische Behandlung der Temporallappenepilepsie. Abstracted in *Zbl. Ges. Neurol. Psychiat.*, 185:1, 1966.
- FARMER, THOMAS W. (Ed.): *Pediatric Neurology*. New York, Harper, 1964.
- FAXÉN, NILS: Le pronostic des convulsions de l'enfance. *Rev. Franc. Pediat.*, 11:665, 1935.
- FEGERSTEN, L.; HAUGE, T.; HENRIKSEN, G. F.; LUNDERVOLD, A., and LOEKEN, A. C.: 28 temporal lobectomies: A preliminary report. *Acta Psychiat. Scand.*, 36:229, 1961.
- FENYES, I.; ZOLTAN, L., and FENYES, G.: Temporal epilepsies with deep-seated epileptogenic foci: Postoperative course. *Arch. Neurol. (Chicago)*, 4:559, 1961.
- FETTERMAN, J., and BARNES, M. R.: Serial studies of the intelligence of patients with epilepsy. *Arch. Neurol. (Chicago)*, 32:797, 1934.
- FOLSOM, ANGELA: Psychological testing in epilepsy, I. Cognitive function. *Epilepsia*, 2:15, 1953.
- FORD, FRANK R.: *Diseases of the Nervous System: In Infancy, Childhood and Adolescence*, 4th ed. Springfield, Illinois, Thomas, 1960.
- FORSTER, FRANCIS M.: Epilepsy. *Traumatic Medicine and Surgery for the Attorney*, 4:481, 1960.
- FOWLER, MALCOLM: Brain damage after febrile convulsions. *Arch. Dis. Child.*, 32:67, 1957.
- FOX, J. TYLOR: The response of epileptic children to mental and educational tests. *Brit. J. Med. Psychol.*, 4:235, 1924.
- FRANTZEN, ESTHER: An analysis of the results of treatment in epileptics under ambulatory supervision. *Epilepsia*, 2:207, 1961.

- FRANTZEN, ESTHER; NYGAARD, ANNE, and WULFF, HENRIK: Febrile convulsions in children: Prognostic studies and attempts at evaluating the effectiveness of a prophylactic anti-epileptic long-term treatment. *Ugeskr. Laeg.*, 126:207, 1964.
- FRENCH, LYLE A.; JOHNSON, DAVID R., and ADKINS, GALEN H.: Cerebral hemispherectomy for intractable seizures . . . A long-term follow-up. *Lancet*, 81:58, (Feb) 1961.
- FRIDERICHSEN, C., and MELCHIOR, J.: Febrile convulsions in children: Their frequency and prognosis. *Acta Paediat.*, 43:307, 1954.
- FRIEDMANN, M.: Ueber die nicht epileptischen Absencen oder kurzen narcoleptischen Anfaelle. *Deutsch. Z. Nervenheilk.*, 30:462, 1906.
- FUKUYAMA, Y.; NAGAHATA, M.; ARIMA, M., and OKADA, Y.: ACTH-Z therapy in flexor spasms in infancy. *Paediat. Univ. Tokyo*, 4:12, 1960.
- FUKUYAMA, YUKIO: Borderlands of childhood epilepsy with special references to febrile convulsions and so-called infantile convulsions. *Proceedings of 11th Annual Meeting of the Japan EEG Society, Tokyo*, 50, (Apr) 1963.
- FUKUYAMA, Y.; ARIMA, M.; NAGAHATA, M., and OKADA, R.: Medical treatment of epilepsies in childhood: A long-term survey of 801 patients. *Epilepsia*, 4:207, 1963.
- GASTAUT, H., *et al.*: A propos du traitement par l'ACTH des encéphalites myocloniques de la première enfance avec dysrythmie majeure (hypsarrhythmia). *Pediatric*, 14:35, 1959.
- GASTAUT, H., *et al.*: H.H.E. syndrome: hemiconvulsions, hemiplegia, epilepsy. *Epilepsia*, 1:418, 1960.
- GASTAUT, H., *et al.*: An electro-clinical study of generalized epileptic seizures of tonic expression. *Epilepsia*, 4:15, 1963.
- GIBBERD, F. B.: The prognosis of petit mal. *Brain*, 89:531, 1966.
- GIBBS, ERNA L.; FLEMING, MARILYN M., and GIBBS, F. A.: Diagnosis and prognosis of hypsarrhythmia and infantile spasms. *Pediatrics*, 13:66, (Jan) 1954.
- GIBBS, ERNA L.; GILLEN, H. W., and GIBBS, F. A.: Disappearance and migration of epileptic foci in childhood. *Amer. J. Dis. Child.* 88:596, 1954.
- GIBBS, ERNA L., and GIBBS, FREDERIC A.: Good prognosis of mid-temporal epilepsy. *Epilepsia*, 1:448, 1960.
- GIBBS, FREDERIC A.: Diagnosis and prognosis of different types of epilepsy. *Cincinnati J. Med.*, 35:409, (Nov) 1954.
- GIBBS, FREDERIC A.; AMADOR, LUIS, and RICH, CATHERINE: Electroencephalographic findings and therapeutic results in surgical treatment of psychomotor epilepsy. In *Temporal Lobe Epilepsy*, Baldwin, M., and Bailey, P. (Eds.). Springfield, Illinois, Thomas, 1958.
- GIBBS, FREDERIC A., and GIBBS, ERNA L.: *Atlas of Electroencephalography*.

- Vol. 1, *Methodology and Controls*, 1st ed. Cambridge, Addison-Wesley, 1941.
- GIBBS, FREDERIC A., and GIBBS, ERNA L.: *Atlas of Electroencephalography*. Vol. 2, *Epilepsy*. Cambridge, Addison-Wesley, 1952.
- GIBBS, FREDERIC A., and GIBBS, ERNA L.: *Atlas of Electroencephalography*. Vol. 3, *Neurological and Psychiatric Disorders*. Reading, Massachusetts, Addison-Wesley, 1964.
- GLIDDON, WILLIAM O.: Gunshot wounds of the head: A review of the after-effects in 500 Canadian pensioners from the Great War, 1914-1918. *J. Canad. Med. Ass.*, 49:373, 1943.
- GLITHERO, E., and SLATER, E.: The schizophrenia-like psychoses of epilepsy, iv. Follow-up record and outcome. *Brit. J. Psychiat.*, 109:134, (Jan) 1963.
- GOLDENSOHN, ELI; O'BRIEN, JOSEPH L., and RANSOHOFF, JOSEPH: Electrical activity of the brain: In patients treated with hemispherectomy or extensive decortication. *Arch. Neurol. (Chicago)* 5:210, (Aug) 1961.
- GOODALL, ROBERT J.: Cerebral hemispherectomy: Present status and clinical indications. *Neurology*, 7:151, 1957.
- GOWERS, W. R.: *Epilepsy and other chronic convulsive diseases: Their causes, symptoms and treatment* (unaltered republication of the work first published by William Wood and Company in 1885), Dover Publications, Inc., New York, 1964.
- GREEN, JOHN; DUISBERG, R. E. H., and MCGRATH, W. B.: Electrocorticography in psychomotor epilepsy. *Electroenceph. Clin. Neurophysiol.*, 3:293, 1951.
- GREEN, JOHN R.; DUISBERG, R. E. H., and MCGRATH, W. B.: Focal epilepsy of psychomotor type: A preliminary report of observations on effects of surgical therapy. *J. Neurosurg.*, 8:157, 1951.
- GREEN, JOHN R., *et al.*: Behavior changes following radical temporal lobe excision in the treatment of focal epilepsy. *Res. Publ. Ass. Res. Nerv. Ment. Dis.*, 36:295, 1958.
- GREEN, JOHN R., and SCHEETZ, DAVID G.: Surgery of epileptogenic lesions of the temporal lobe. *Arch. Neurol. (Chicago)* 10:135, (Feb) 1964.
- GREENSTEIN, LOUIS: Epilepsy and the electroencephalogram. *J. Mount Sinai Hosp. NY*, 19:683, 1953.
- GRINKER, ROY R.; BUCY, PAUL C., and SAIIS, ADOLPH L.: *Neurology*, 5th ed. Springfield, Illinois, Thomas, 1960.
- GROSZ, WOLFGANG: Ueber den Ausgang der genuinen Epilepsie: (auf Grund kataamnestischer Erhebungen). *Archiv. Psychiat.*, 90:765, 1930.
- GRUILE, HANS W.: Ueber die Fortschritte in der Erkenntnis der Epilepsie in den Jahren 1910-1920 und ueber das Wesen dieser Krankheit. *Zbl. Ges. Neurol. Psychiat.*, 34:1, 1924.
- GUILLAUME, JEAN; MAZARS, G., and MAZARS, Y.: Étude statistique des ré-

- sultats éloignés de la chirurgie de l'épilepsie. *Presse Med.*, 61:881, (June) 1953.
- GUTTMANN, IRENE: Beitrag zur Epilepsiestatistik. *Z. Ges. Neurol. Psychiat.*, 118:500, 1929.
- HABERMAAS: Ueber die Prognose der Epilepsie. *Z. Psychiat.*, 58:243, 1901.
- HALSTEAD, H.: Abilities and behaviour of epileptic children. *J. Ment. Sci.*, 102:28, 1957.
- HARRIS, R., and TIZARD, J. P. M.: The electroencephalogram in neonatal convulsions. *J. Pediat.* 57:501, (Oct) 1960.
- HARRIS, RUTH: Some EEG observations in children with infantile spasms treated with ACTH. *Arch. Dis. Child.*, 39:564, 1964.
- HEDENSTROEM, INGE, and SCHORSCH, GERHARD: Klinische und hirnelektrische Befunde bei 120 anfallsfrei gewordenen Epileptikern. *Archiv. Psychiat. Nervenkr.*, 198:17, 1958.
- HEDENSTROEM, INGE, and SCHORSCH, GERHARD: Ueber therapieresistente Epileptiker. *Archiv. Psychiat. Nervenkr.*, 204:579, 1963.
- HENDERSON, P.: Epilepsy in school children. *Brit. J. Prev. Soc. Med.*, 7:9, 1953.
- HENRIKSEN, P. B.; JUUL-JENSEN, P., and LUND, M.: The Mortality of Epileptics, A Preliminary Report. *Social Studies in Epilepsy, No. 5*, International Bureau for Epilepsy, London, 1966, p. 5.
- HERLITZ, GILLIS: Studien ueber die sog. initialen Feiberkraempfe bei Kindern, with an English summary. *Acta Paediat.*, 29:1, 1941.
- HERTOFT, P.: The clinical, electroencephalographic and social prognosis in petit mal epilepsy. *Epilepsia*, 4:298, 1963.
- HESS, RUDOLF: Verlaufsuntersuchungen ueber Anfaelle und EEG bei kindlichen Epilepsien. *Archiv. Psychiat. Nervenkr.*, 197:568, 1958.
- HILL, DENIS: Indications and contra-indications to temporal lobectomy. *Proc. Roy. Soc. Med. [Neurol.]*, 51:610, (Feb) 1958.
- HILL, DENIS: The E.E.G. in psychiatry. In *Electroencephalography*, Hill, D., and Parr, G. (Eds.). New York, Macmillan, 1963.
- HIMLER, LEONARD E., and RAPHAEL, THEOPHILE: A follow-up study on 93 college students with epilepsy. *Amer. J. Psychiat.*, 101:760, 1945.
- HIPPOCRATES: Hippocratic writings, trans. by Francis Adams. In *Great Books of the Western World*, vol. 10 Chicago, Encyclopedia Britannica, 1952.
- HOFER, PAUL F. A.; HOFF, HANS H., and PLUVINAGE, ROGER I. L.: The effect of anticonvulsant therapy on the electroencephalogram of patients with idiopathic epilepsy. *Trans. Amer. Neurol. Ass.*, 185, 1947.
- HOLOWACH, JEAN; THURSTON, DONALD L., and O'LEARY, JAMES L.: Petit mal epilepsy. *Pediatrics*, 30:893, (Dec) 1962.
- HORSLEY, VICTOR: Surgery of the central nervous system. *Brit. Med. J.*, 2:1286, (Dec) 1890.

- HORSTMANN, WINFRED, and SCHINNERLING, WALTER: Zur Prognose der sog. Fieberkrampe. *Mscr. Kinderheilk.*, 111:52, 1963.
- HRBEK, ANTONÍN: Fieberkrampe im Kindesalter. Febrile fits in childhood—Convulsions hyperthermiques chez l'enfant. *Ann. Paediat.*, (Basel), 188:162, 1957.
- HUNTER, RICHARD A.: Status epilepticus: History, incidence and problems. *Epilepsia*, 1:162, 1959-60.
- HYLLESTED, KAY, and PAKKENBERG, H.: Prognosis in epilepsy of late onset. *Neurology*, 13:641, 1963.
- ILLINGWORTH, R. S.: Sudden mental deterioration with convulsions in infancy. *Arch. Dis. Child.*, 30:529, 1955.
- ISLER, WERNER, and HESS, RUDOLF: Verlaufsuntersuchungen ueber Anfaelle und EEG bei fokalen Epilepsien im Kindesalter. *Arch. Psychiat. Nervenkr.*, 200:257, 1960.
- JACKSON, JOHN HUGHLINGS, *Selected Writings of*: Volume One on Epilepsy and Epileptiform Convulsions (Edited by James Taylor), Staples Press, London, 1958.
- JANZ, D.: Die klinische Stellung der Pyknolepsie. *Deutsch. Med. Wschr.*, 10:1392, (Sept) 1955.
- JANZ, D.: The grand mal epilepsies and the sleep-waking cycle. *Epilepsia*, 3:69, 1962.
- JASPER, H.; PERTUISSET, B., and FLANIGIN, H.: EEG and cortical electrograms in patients with temporal lobe seizures. *Arch. Neurol. Psychiat.*, 65:272, 1951.
- JASPER, H. H.; ARFEL-CAPDEVILLE, G., and RASMUSSEN, T.: Evaluation of EEG and cortical electrographic studies for prognosis of seizures following surgical excision of epileptogenic lesions. *Epilepsia*, 2:130, 1961.
- JEAVONS, P. M., and BOWER, B. D.: The natural history of infantile spasms. *Arch. Dis. Child.*, 36:17, 1961.
- JENKINS, C. DAVID: The relation of EEG slowing to selected indices of intellectual impairment. *J. Nerv. Ment. Dis.*, 135:162, 1962.
- JENNETT, W. BRYAN, and LEWIN, WALPOLE: Traumatic epilepsy after closed head injuries. *J. Neurol. Neurosurg. Psychiat.*, 23:295, 1960.
- JENNETT, W. BRYAN: Predicting epilepsy after blunt head injury. *Brit. Med. J.*, 1:1215, (May) 1965.
- JENNINGS, C. G.: Febrile convulsions in childhood, review of their significance. *J. Mich. Med. Soc.*, 53:272, (Mar.) 1954.
- JOEDICKE, P.: Ueber die aetiologischen Verhaeltnisse, Lebensdauer, allgemeine Sterblichkeit, Todesursachen und Sektionsbefunde bei Epileptikern. Abstracted in *Zbl. Ges. Neurol. Psychiat.*, 8:194, 1914.
- JØRGENSEN, R. SØGAARD: Follow-up examination of epileptic patients. *Acta Psychiat. Scand.*, 36:225, 1961.
- JUUL-JENSEN, PALLE: *Epilepsy, A Clinical and Social Analysis of 1020 Adult Patients with Epileptic Seizures*. Copenhagen, Munksgaard, 1963.

- JUUL-JENSEN, PALLE: Frequency of recurrence after discontinuance of anti-convulsant therapy in patients with epileptic seizures. *Epilepsia*, 5:352, 1964.
- KEITH, HADDOW M.: The treatment of epilepsy in children. *J. Canad. Med. Ass.*, 37:485, 1937.
- KEITH, HADDOW M.: Results of treatment of recurring convulsive attacks of epilepsy. *Amer. J. Dis. Child.*, 74:140, 1947.
- KEITH, HADDOW M.; EWERT, JOSEPHINE C.; GREEN, MEREDITH, W., and GAGE, ROBERT P.: Mental status of children with convulsive disorders. *Neurology*, 5:419, 1955.
- KEITH, HADDOW M.: *Convulsive Disorders in Children With Reference to Treatment with Ketogenic Diet*. Boston, Little, 1963.
- KEITH, HADDOW M.: Convulsions in children under three years of age: A study of prognosis. *Mayo Clin. Proc.*, 39:895, (Dec.) 1964.
- KENNEDY, W. A., and HILL, DENIS: The surgical prognostic significance of the electroencephalographic prediction of Ammon's horn sclerosis in epileptics. *J. Neurol., Neurosurg. Psychiat.*, 21:24, 1958.
- KJØRBOE, ERIK; LUND, MOGENS, and POULSEN, ANNEISE: The prognosis of epilepsy of short duration in adults, with special regard to the risk in traffic and industry. *Ugeskr. Laeg.*, 2:10, 1958.
- KJØRBOE, ERIK: The prognosis of epilepsy. *Acta Psychiat. Scand.*, 36:166, 1961.
- KIRSTEIN, LENNART: A contribution to the knowledge of the prognosis of epilepsy. *Acta Med. Scand.*, 112:515, 1942.
- KOCH, FR., and GRUETZNER, A.: Zur ACTH-Behandlung der Blitz-Nick-Salaam-Krämpfe des Säuglings. *München. Med. Wschr.*, 102:36, 1960.
- KROHN, W.: Causes of death among epileptics. *Epilepsia*, 4:315, 1963.
- KUGELMASS, I. NEWTON; POULL, LOUISE E., and RUDNICK, J.: Mental growth of epileptic children. *Amer. J. Dis. Child.*, 55:295, 1938.
- KUHLO, WOLFGANG: Katamnestic Untersuchungen zur Pyknolepsie. *Arch. Psychiat. Nervenkr.*, 207:254, 1965.
- LAMONTAGNE, A., and FISCHGOLD, H.: Action des anticonvulsivants étude longitudinale d'un groupe de jeunes épileptiques. *Thérapie*, 20:1354, 1965.
- LANDOLT, H.: Die Bedeutung der Elektroencephalographie fuer die Behandlung der Epilepsie. *Nervenarzt*, 28:170, (Jan) 1957.
- LEES, P., and LAYESEDGE, L. A.: The prognosis of "petit mal" and minor epilepsy. *Lancet*, 797 (Oct) 1962.
- LEHTINEN, LAURI O.: Prognosis of epilepsy: A preliminary report. *Acta Neurol. Scand.*, 41:517, 1965.
- LEMP, R.: Die cerebralen Anfallskrankheiten im Kindesalter. In Schulte, Walter (Ed.): *Epilepsie und ihre Randgebiete in Klinik und Praxis*. Germany, J. F. Lehmanns Verlag Muenchen, 1964.
- LENORMANT, CH.: Quelques considérations sur l'épilepsie consécutive aux

- traumatismes du crâne et son traitement. Frequence de l'épilepsie traumatique et conditions qui la déterminent. Données anatomo-physiologiques sur lesquelles se base l'intervention chirurgicale. Technique opératoire. Résultats. *Zbl. Ges. Neurol. Psychiat.*, 28:501, 1922.
- LENNOX, MARGARET A.: Febrile convulsions in childhood, a clinical and electroencephalographic study. *Amer. J. Dis. Child.*, 78:868, 1949.
- LENNOX, MARGARET A.: Personal communication, 1967.
- LENNOX, WILLIAM GORDON: *Epilepsy and Related Disorders*, Vols. 1 and 2. Boston, Little, 1960.
- LENNOX, WILLIAM G., and COLLINS, A. L.: Intelligence of normal and epileptic twins. *Amer. J. Psychiat.*, 101:764, 1945.
- LIVINGSTON, S.; BRIDGE, E., and KAJDI, L.: Febrile convulsions: A clinical study with special reference to heredity and prognosis. *J. Pediat.*, 31:509, 1947.
- LIVINGSTON, SAMUEL: *The Diagnosis and Treatment of Convulsive Disorders in Children*. Springfield, Illinois, Thomas, 1954.
- LIVINGSTON, SAMUEL, and PETERSEN, DON: Primidone (Mysoline) in the treatment of epilepsy. *New Eng. J. Med.*, 254:327, (Feb) 1956.
- LIVINGSTON, S.; EISNER, V., and PAULI, LYDIA: Minor motor epilepsy: Diagnosis, treatment and prognosis. *Pediatrics*, 21:916, (June) 1958.
- LIVINGSTON, SAMUEL: Living with epileptic seizures. *Pediatrics*, 59:128, 1961.
- LIVINGSTON, SAMUEL: *Living with Epileptic Seizures*. Springfield, Illinois, Thomas, 1963.
- LIVINGSTON, S.; TORRES, I.; PAULI, L. L., and RIDER, R. V.: Petit mal epilepsy: Results of a prolonged follow-up study of 117 patients. *JAMA*, 194:227, (Oct) 1965.
- LORGÉ, M.: Epilepsie und Lebensschicksal: Ergebnisse katamnestischer Untersuchungen, *Psychiat. Neurol. (Basel)*, 147:360, 1964.
- LUND, M.: Diseased, Disabled and Elderly Persons in Road Traffic: The Prognosis of Epilepsy of Short Duration in Adults. *The Second Congress of the International Association for Accident and Traffic Medicine*, Section II:B 1, Stockholm, 1966.
- LUNDVOLD, ARNE, and JABBOUR, J. T.: A correlation of clinical, electroencephalographic, and roentgenographic findings in children with epilepsy. *J. Pediat.*, 60:220, 1962.
- LUNDVOLD, ARNE: A prognostic evaluation of clinical, electroencephalographic and roentgenographic findings in children with epilepsy. *Epilepsia*, 5:33, 1964.
- MAKI, Y.: Prognosis of early post-traumatic epilepsy. *Excerpta Med.*, [VIII], 19:637, 1966.
- MARSHALL, CURTIS, and WALKER, EARL: The value of electroencephalography in the prognostication and prognosis of post-traumatic epilepsy. *Epilepsia*, 2:138, 1961.

- MASQUIN, A., and COURJON, J.: Prognostic factors in posttraumatic epilepsy. *Epilepsia*, 4:285, 1963.
- MATERA, RAUL, and CASTRO, MANUEL, M.: Hemisferectomias, *Sem. Med. (B Air)*, 123:1995, (Dec) 1963.
- MATTHES, A.: Die Blitz-, Nick- und Salaamkrampe und ihre Prognose *Mtschr. Kinderheilk.*, 102:68, 1954.
- MATTHIOLIUS: Ueber Schadeloperationen bei Epilepsie. *Deutsch. Z. Chirurgie*, 52:1, 1899.
- McKISSOCK, WYLIE: Personal communication, (Feb) 1967.
- MELIN, KARL-AXEL: Zur Prognose der Infektkrampe. *Mtschr. Kinderheilk.*, 102:62, 1954.
- MERRITT, H. HOUSTON: Paroxysmal disorders, convulsive disorders, epilepsy. In *Textbook of Neurology*, 3rd ed. Philadelphia, Lea and F. 1963.
- MESSINA, CORRADO, and RAGONESE, GIUSEPPE: Correlazioni fra risultati clinici ed elettroencefalografici ottenuti in corso di terapia anticomiziale. *Riv. Neurol.*, 35:411, 1965.
- MEYERS, RUSSELL: The surgical treatment of "focal" epilepsy: An inquiry into current premises, their implementation and the criteria employed in reporting results. *Epilepsia*, 3:9, 1954.
- MICHAUX, M. LÉON: Les graves conséquences des pronostics abusifs d'incurabilité chez les enfants et les adolescents mentalement inadaptés. *Bull. Acad. Nat. Med. (Paris)*, 148:129, 1964.
- MILLER, JAMES G.: The organization of life. *Perspect. Biol. Med.*, 9:107, 1965.
- MILLICHAP, J. G.; ALEDORT, L. M., and MADSEN, J. A.: A critical evaluation of therapy of febrile seizures. *J. Pediat.*, 56:364, (Mar) 1960.
- MILLICHAP, J. GORDON; MADSEN, JACK A., and ALEDORT, LOUIS M.: Studies in febrile seizures, V. Clinical and electroencephalographic study in unselected patients. *Neurology*, 10:643, (July) 1960.
- MONRAD, S.: *Epilepsie im Kindesalter*. Abstracted in *Zbl. Ges. Neurol. Psychiat.*, 33:122, 1923.
- MONRAD, S.: Die Prognose und die Behandlung der Kinder-epilepsie. *Acta Paediat.*, 12:231, 1932.
- MORRIS, ARTHUR A.: The surgical treatment of psychomotor epilepsy. *Med. Ann. DC*, 19:121, 1950.
- MORRIS, ARTHUR A.: Temporal lobectomy with removal of uncus, hippocampus, and amygdala: Results for psychomotor epilepsy three to nine years after operation. *Arch. Neurol. Psychiat. (Chicago)*, 76:479, 1956.
- MUNSON, J. F.: Death in epilepsy. *Med. Rec.*, 77:58, (Jan) 1910.
- MUSKENS, L. J. J.: *Epilepsy, Comparative Pathogenesis, Symptoms, Treatment*. London, Baillière, Tindall and Cox, 1928.

- NORTHFIELD, D. W. C.: Discussion on surgery of temporal lobe epilepsy. *Proc. Roy. Soc. Med., [Neurol.]*, 51:607, (Feb) 1958.
- OBRADOR, S., and LARRAMENDI, L. M. H.: Clinical and electroencephalographic follow-up of four epileptics treated by cerebral hemispherectomy. *Electroenceph. Clin. Neurophysiol.*, 3:63, 1953.
- OPPENHEIM, H.: *Lehrbuch der Nervenkrankheiten fuer Aerzte und Studierende*, Berlin, 1908.
- OSTMANN: Gesammelte Notizen ueber unsere Epileptiker, ein Beitrag zur Kenntnis der Epilepsie. *Zbl. Ges. Neurol. Psychiat.*, 52:725, 1929.
- PAAL, G.: Katamnestic Untersuchungen und EEG bei Pyknolepsie. *Arch. Psychiat. Nervenkr.*, 196:48, 1957.
- PACHE, HANS-DIETRICH: Die Klinik der kindlichen Krampfleiden. *Mtschr. Kinderheilk.*, 102:42, 1954.
- PAILLAS, J. E.: Aspects cliniques de l'épilepsie temporale. In *Temporal Lobe Epilepsy*, Baldwin, M., and Bailey, P. (Eds.) Springfield, Illinois, Thomas, 1958.
- PASKIND, HARRY A., and BROWN, MEYER: Constitutional differences between deteriorated and non-deteriorated patients with epilepsy: II. Anthropometric measurements. *Amer. J. Psychiat.*, 95:901, 1939.
- PASSOUANT, P., and CADILHAC, J.: EEG and clinical study of epilepsy during maturation in man. *Epilepsia*, 3:14, 1962.
- PATRICK, HUGH T., and LEVY, DAVID M.: Early convulsions in epileptics and in others. *JAMA*, 82:375, (Feb) 1924.
- PATTERSON, HAROLD A., and FONNER, DELMA: Some observations on the intelligence quotient in epileptics, a preliminary report. *Psychiat. Quart.*, 31:542, (Oct) 1928.
- PAULI, L.; O'NEIL, R.; YBANEZ, M., and LIVINGSTON, S.: Minor motor epilepsy, treatment with Corticotropin (ACTH) and steroid therapy. *JAMA*, 174:152, (Nov) 1960.
- PEIFFER, JUERGEN: *Morphologische Aspekte der Epilepsien, Pathogenetische, Pathologisch-Anatomische und Klinische Probleme der Epilepsien*. Berlin, Springer-Verlag, 1963.
- PENFIELD, W., and STEELMAN, H.: The treatment of focal epilepsy by cortical excision. *Am. Surg.*, 126:740, 1947.
- PENFIELD, WILDER, and FLANIGIN, H.: Surgical therapy of temporal lobe seizures. *Arch. Neurol. Psychiat. (Chicago)* 64:491, 1950.
- PENFIELD, WILDER, and JASPER, HERBERT: *Epilepsy and the Functional Anatomy of the Human Brain*. Boston, Little, 1954.
- PENFIELD, WILDER, and PAINE, KENNETH: Results of surgical therapy for focal epileptic seizures. *Canad. Med. Ass. J.*, 78:515, 1955.
- PERKINS, ROY F., and LAUFER, MAURICE W.: Follow-up study on eighty cases of convulsive disorder. *J. Nerv. Ment. Dis.*, 106:159, 1947.
- PETERMAN, M. G.: Febrile convulsions in children. *JAMA*, 143:728, (June) 1950.

- PETERMAN, M. G.: Febrile convulsions. *J. Pediat.*, 41:536, (Nov) 1952.
- PETIT-DUTAILLIS, D.; CHRISTOPHE, J.; PERTUISSET, B., and DREYFUS-BRISAC, C.: Indications et résultats de l'excision corticale dans l'épilepsie temporaire (d'après 20 cas opérés). *Sem. Hop. Paris*, 29:3838, 1953.
- PHILLIPS, GILBERT: Traumatic epilepsy after closed head injury. *J. Neurol., Neurosurg. Psychiat.*, 17:1, 1954.
- PICAZA, J. A., and GUMA, J.: Experience with the surgical treatment of psychomotor epilepsy. *Arch. Neurol. and Psychiat. (Chicago)* 75:57, 1956.
- PIGOTT, ALBERT W.; WEINGROW, S. M., and FITCH, THOMAS S. P.: Convulsions in the chronic nervous diseases of infancy and childhood, a review of 1,660 cases. *Arch. Pediat.*, 56:703 and 781, 1939; 57:12 and 92, 1940.
- POHLISCH, KURT: Zur Frage der Pyknolepsie, gehäufte kleine Anfälle der Kinder. *Arch. Psychiat. Nervenkr.*, 67:424, 1923.
- POND, D. A.; BIDWELL, B. H., and STEIN, L.: A survey of epilepsy in fourteen general practices, I. Demographic and medical data. *Psychiat., Neurol., Neurochir.*, 63:217, 1960.
- POND, D. A., and BIDWELL, B. H.: A survey of epilepsy in fourteen general practices, II. Social and psychological aspects. *Epilepsia*, 1:285, 1960.
- PRICHARD, J. S., and MCGREAL, D. A.: Febrile convulsions. *Med. Clin. N. Amer.*, Mar. 1958, p. 379.
- PROBST, CHARLES: Ueber den Verlauf von hirnelektrisch stummen Epilepsien. *Schweiz. Arch. Neurol. Psychiat.*, 85:357, 1960.
- PUSSEP, L.: Die chirurgische Behandlung der Epilepsie (nach zwanzigjährigen eigenen Beobachtungen). *Klin. Wschr.* 1:2142, (Oct) 1922.
- QUADFASSEL, ANGELA FOLSOM, and PRUYSER, PAUL W.: Cognitive deficit in patients with psychomotor epilepsy. *Epilepsia*, 4:80, 1955.
- RANHEIM, B.; DIETRICHSON, P., and WILLIAMSEN, R.: Prognosis in epilepsy, II. Further report of an investigation. *Acta Neurol. Scand.*, (Suppl. 41). 13:497, 1965.
- RANSOHOFF, J., and CARTER, S.: Hemispherectomy in the treatment of convulsive seizures associated with infantile hemiplegia. *Proc. Res. Nerv. Ment. Dis.*, 34:176, 1954.
- RASMUSSEN, T., and JASPER, H.: Temporal lobe epilepsy, indication for operation and surgical technique. In *Temporal Lobe Epilepsy*, Baldwin, M., and Bailey, P. (Eds.). Springfield, Illinois, Thomas, 1958.
- RASMUSSEN, THEODORE, and BRANCH, CHARLES: Temporal lobe epilepsy indications for and results of surgical therapy. *Postgrad. Med.*, 31:9 (Jan) 1962.
- RASMUSSEN, T.: Surgical therapy of frontal lobe epilepsy. *Epilepsia*, 4:181 1963.
- RAWLING, L. BATHE: The remote effects of gunshot wounds of the head. *Brit. J. Surg.*, 10:93, 1922-1923.

- REDLICH, EMIL: Zur Pathologie der Epilepsie nach Schaedelschussverletzungen. *Z. Ges. Neurol. Psychiat.*, 48:8, 1919.
- RUSSELL, R. W., and WHITTY, C. W. M.: Studies in traumatic epilepsy, I. Factors influencing the incidence of epilepsy after brain wounds. *J. Neurol., Neurosurg. Psychiat.*, 15:93, 1952.
- RODIN, E. A.: An electroencephalographic syndrome which correlates with severe disruption of the personality structure. *Neurology*, 7:724, 1957.
- RODIN, E. A.: A system of coding EEG reports for use by electronic computers. *Electroenceph. Clin. Neurophysiol.*, 13:795, 1961.
- RODIN, E. A.; BECKETT, P. G., and SOKOLOV, S. L.: A coding system for patients with convulsive disorders and children with behavior disturbances. *IRE Transactions on Bio-Medical Electronics*, 1:61, (Jan.) 1962.
- RODIN, E. A.: Some relationships of induced seizure patterns to clinical findings in epileptic patients. *Epilepsia*, 5:21, 1964.
- RODIN, E. A., LIN, Y., GONZALEZ, S., and DENNERLL, R. D.: The application of computer techniques to clinical research in convulsive disorders. *IEEE Trans. Biomed. Engineering*, 11:19, (Jan/April) 1964.
- RODIN, E. A., *et al.*: The electroencephalogram in acute fatal head injuries. *J. Neurosurg.*, 23:329, 1965.
- RODIN, E. A., and GONZALEZ, S.: Hereditary components in epileptic patients, EEG family studies. *JAMA*, 198:221, (Oct.) 1966.
- ROMANO, J., and ENGEL, G. L.: Delirium, I. Electroencephalographic data. *Arch. Neurol. Psychiat.*, 51:356, 1944.
- ROSAS, FREDERICO SAL Y.: Convulsions in childhood and epilepsy. *J. Nerv. Ment. Dis.*, 121: 236, 1955.
- ROSENTHAL, CURT: II. Die gehaeuften kleinen Anfaelle des Kindesalters (Pyknolepsie). *Ergebn. Inn. Med. Kinderheilk.*, 48:77, 1935.
- SARGENT, PERCY: Some observations on epilepsy. *Brain*, 44:312, 1921.
- SAUER, H.: Ueber gehaeufte kleine Anfaelle bei Kindern (Pyknolepsie). *Mschr. Psychiat. Neurol.*, 40:276, 1916.
- SCHMIDT, G.: Die Blitz-Nick-Salaamkraempfe und ihre Prognose nach Hormonbehandlung. *Mschr. Kinderheilk.*, 112:6, 1964.
- SCHWADE, EDWARD, and OTTO, OWEN: Mortality in epilepsy. *JAMA*, 156: 1526, (Dec.) 1954.
- SELBACH, HELMUT: 12. Die cerebralen Anfallsleiden: Genuine Epilepsie, symptomatische Hirnkrampfe und die Narkolepsie. In *Handbuch der inneren Medizin*, 4th ed., 3:1082, Berlin, Germany, Springer, 1953.
- SERAFETINIDES, E. A., and FALCONER, M. A.: The effects of temporal lobectomy in epileptic patients with psychosis. *J. Ment. Sci.*, 108:584, 1962.
- SERAFETINIDES, E. A.; DOMINIAN, J., and DEWHURST, M.: A follow-up study of late onset epilepsy, I. Neurological findings, II. Psychiatric and social findings. *Brit. Med. J.*, 1:428, (Feb.) 1963.
- SHANKS, ROBERT A.: Convulsions in childhood and their relation to epilepsy. *Glasgow Med. J.*, 32:257, (Sept.) 1951.

- SIMMEL, MARIANNE L., and COUNTS, SARAH: Clinical and psychological results of anterior temporal lobectomy in patients with psychomotor epilepsy. In *Temporal Lobe Epilepsy*, Baldwin, M., and Bailey, P. (Eds.). Springfield, Illinois, Thomas, 1958.
- SLATER, ELIOT, and BEARD, A. W.: The schizophrenia-like psychoses of epilepsy, i. Psychiatric aspects, v. Discussion and conclusions. *Brit. J. Psychiat.*, 109:95, 1963.
- SLATER, ELIOT, and GLITHERO, ERIC: The schizophrenia-like psychoses of epilepsy, iii. Genetical aspects. *Brit. J. Psychiat.*, 109:95, 1963.
- SOREL, L., and DUSAUCY-BAULOYE, A.: A propos de 21 cas d'hypsarhythmia de Gibbs, son traitement spectaculaire par l'ACTH. *Acta Neurol. Belg.*, 58:130, (Feb.) 1958.
- SPIEGEL, E. A.; WYCIS, H. T., and BAIRD, H. W., III: Long-range effects of electropallidoansotomy in extrapyramidal and convulsive disorders. *Neurology*, 8:734, 1958.
- SPIEGEL, E. A.; WYCIS, H. T., and BAIRD, H. W., III: Pallidotomy and pallidoamygdalotomy in certain types of convulsive disorders. *Arch. Neurol. Psychiat. (Chicago)*, 80:714, (Dec.) 1958.
- STAMPS, F. W., et al.: Treatment of hypsarhythmia with ACTH. *JAMA*, 171:408, (Sept.) 1959.
- STEINSIEK, HANS-DIETRICH: Ueber Todesursachen und Lebensdauer bei genuiner Epilepsie. *Arch. Psychiat. Nervenkr.*, 183:469, 1950.
- STEVENSON, W. E.: Epilepsy and gunshot wounds of the head. *Brain*, 54: 214, 1931.
- STOLECKE, H., and PACHE, H.-D.: Zur Behandlung von BNS-Krampfen mit ACTH. *Msehr. Kinderheilk.*, 110:105, 1962.
- STOLLER, ALAN: Slowing of the alpha-rhythm of the electroencephalogram and its association with mental deterioration and epilepsy. *J. Ment. Sci.* 95:972, (Oct) 1949.
- STONE, THEODORE T., and ARIEFF, ALEX J.: Remissions in epileptic patients treated with sodium bromide in an outpatient clinic. *Arch. Neurol. Psychiat. (Chicago)*, 43:299, 1940.
- STROBOS, ROBERT R. J.: Prognosis in convulsive disorders. *Arch. Neurol. (Chicago)*, 1:216, 1959.
- SULLIVAN, ELLEN B., and GALLAGAN, LAWRENCE: On intelligence of epileptic children. *Genet. Psychol. Monog.*, 17:309, (Oct.) 1935.
- SYMONDS, C. P.: Traumatic epilepsy. *Lancet*, 2:1217, (Nov.) 1935.
- TENNY, JOHN W.: Epileptic children in Detroit's special school program, a study. *Exceptional Child.*, 21:162, (Feb.) 1955.
- THOM, DOUGLAS A.: Convulsions of early life and their relation to the chronic convulsive disorders and mental defect. *Amer. J. Psychiat.*, 98: 574, (Jan.) 1942.
- THOMASKE, G.; WOHINGER, E., and SCHNEIDER, J.: 1. Operative Behand-

- lung der Temporallappenepilepsie, Nachuntersuchung chirurgisch behandelte Epileptiker. *Zbl. Ges. Neurol. Psychiat.*, 185:3, 1966.
- TIBBLES, J. A. R., and PRICHARD, J. S.: The prognostic value of the electroencephalogram in neonatal convulsions. *J. Pediat.*, 35:778, (May) 1965.
- TILMANN: Die chirurgische Behandlung der traumatischen Epilepsie. *Arch. Klin. Chirurgie*, 92:496, 1910.
- TOMAN, JAMES E. P.: The neuropharmacology of antiepileptics. *Electroenceph. Clin. Neurophysiol.*, 1:33, 1949.
- TROJABORG, W., and PLUM, P.: Treatment of "hypsarrhythmia" with ACTH. *Acta Paediat.*, 49:572, (Sept.) 1960.
- TROLLE, ELLI: Drug therapy of epilepsy. *Acta Psychiat. Scand. (Suppl. 150)*, 36:187, 1961.
- TURNER, WILLIAM ALDREN: *Epilepsy, A Study of the Idiopathic Disease*. London, Macmillan, 1907.
- TURNER, ERIC: A new approach to unilateral and bilateral lobotomies for psychomotor epilepsy. *J. Neurol. Neurosurg. Psychiat.*, 26:285, 1963.
- ULRICH, J.: Verlaufuntersuchungen bei sensiblen und motorischen fokalen Epilepsien. *Schweiz. Med. Wschr.*, 91:1226, 1961.
- WAGSTAFFE, W. W.: The incidence of traumatic epilepsy after gunshot wound of the head. *Lancet*, 2:861, (Oct.) 1928.
- WAKANA, A., et al.: Relationship between pre-therapy EEG and the prognosis of epileptics. *Excerpta Med. [VIII]*, 18:14, 1965.
- WALKER, A. EARL: Prognosis in post-traumatic epilepsy, a ten-year follow-up of craniocerebral injuries of World War II. *JAMA*, 164:1636, (Aug.) 1957.
- WALKER, A. EARL, and JABLON, SEYMOUR: A follow-up of head-injured men of World War II. *J. Neurosurg.*, 26:600, 1959.
- WALKER, A. EARL, and JABLON, SEYMOUR: A follow-up study of head wounds in World War II. *VA Medical Monograph*, U. S. Government Printing Office, Washington 25, D.C., 1961.
- WALKER, A. EARL: Post-traumatic epilepsy. *World Neurol.*, 3:185, (Mar.) 1962.
- WATSON, C. WESLEY: The incidence of epilepsy following cranio-cerebral injury. *Res. Publ. Ass. Res. Nerv. Ment. Dis.*, 26:516, 1947.
- WATSON, C. WESLEY: Incidence of epilepsy following craniocerebral injury, II. Three-year follow-up study. *Arch. Neurol. Psychiat. (Chicago)*, 68:831, 1952.
- WEST, W. J.: On a peculiar form of infantile convulsions (a letter to the editor). *Lancet*, January 26, 1841, p. 724.
- WILKINS, LAWSON: Epilepsy in childhood, I. A statistical study of clinical types. *J. Pediat.*, 10:317, 1937.
- WILLIAMS, DENIS: The electroencephalogram in traumatic epilepsy. *J. Neurol., Neurosurg. Psychiat.*, 7:103, (Oct.) 1944.

- WOODCOCK, S., and COSGROVE, J. B. R.: Epilepsy after the age of 50, A five-year follow-up study. *J. Neurol.*, 14:34, (Jan.) 1964.
- WORINGER, E.; BAUMGARTNER, J.; THOMALSKE, G., and SCHNEIDER, J.: Résultant du traitement neuro-chirurgical de 22 de nos épilepsies à foyer électroencéphalographique temporal. *Neurochirurgie*, 1:385, 1955.
- WORLD HEALTH ORGANIZATION: *Juvenile Epilepsy, Report of a Study Group*. Report Series No. 130, Geneva (Switzerland), 1957.
- YACORZYNSKI, C. K., and ARIEFF, ALEX J.: Absence of deterioration in patients with non-organic epilepsy with especial reference to bromide therapy. *J. Nerv. Ment. Dis.*, 95:687, 1942.
- YAHN, M.; SCIARRA, D.; CARTER, S., and MERRITT, H.: Evaluation of standard anticonvulsant therapy in 319 patients. *JAMA*, 150:663, 1952.
- YAHN, MELVIN D., and MERRITT, H. HOUSTON: Current status of the drug therapy of epileptic seizures. *JAMA*, 161:333, (May) 1956.
- YAHN, MELVIN D., and MERRITT, H. HOUSTON: The drug therapy of convulsive seizures. *Int. J. Neurol.*, 1:76, (Dec.) 1959.
- YANNET, HERMAN (Chairman); DEAMER, WILLIAM C., and BARBA, PHILIP S.: Round table discussion, convulsive disorders in children. *J. Pediat.* Nov., 1949, p. 677.
- ZELLWEGER, HANS: *Krämpfe im Kindesalter, I. Teil*. Basel, Schwabe, 1948.
- ZELLWEGER, HANS: Convulsions in childhood. *Krämpfe im Kindesalter*. Convulsions chez l'enfant. *Ann. Paediat. (Basel)*, 190:257, (May) 1958.
- ZENKER, C.; GROH, C., and ROTH, G.: Probleme und Erfahrungen beim Absetzen antikonvulsiver Therapie. *Z. Kinderheilk.*, 2:152, 1957.
- ZIMMERMAN, F. T.; BURGEMEISTER, BESSIE B., and PUTNAM, T. J.: Intellectual and emotional makeup of the epileptic. *Arch. Neurol. Psychiat. (Chicago)*, 65:545, (May) 1951.
- ZIMMERMAN, H. M.: The histopathology of convulsive disorders in children. *J. Pediat.*, 13:859, 1938.

AUTHOR INDEX

- Aass, F., 41
 Abbott, J. A., 16
 Adie, W. J., 54, 59
 Ajmone-Marsan, C., 117t, 120
 Alajouanine, Th., 89t, 99t
 Astroem, C. H., 4-8, 4t, 7t, 10, 13-14, 162, 169
 Annmann, —, 159, 170t
 Arieff, A. J., 4-5, 11-14, 145-146, 162
 Arnold, O. H., 132
 Ascroft, P. B., 89-99, 89t, 90ts, 91t, 93t, 99t

 Bailey, P., 117-118, 117t, 120
 Baldwin, M., 117t, 120
 Bamberger, P., 133
 Barnes, M. R., 143
 Barrow, R. L., 3
 Bauman, H., 88-90, 89t, 90ts, 92-94, 97, 97t, 99t
 Bergamini, L., 67, 71t
 Bergemann, E., 32
 Bloom, D., 120
 Bower, B. D., 75, 77, 79-80, 84-85
 Brain, W. R., 9-10, 14
 Branch, C., 117-120, 117t
 Bray, P. F., 80-81
 Breg, W. R., 23
 Bridge, E. M., 5-6, 6t, 24t, 27, 30, 32-33, 60-61, 133, 138, 160, 168
 Brown, M., 148
 Brun, R., 89t
 Burke, J. B., 31-32, 168-169

 Cadillac, J., 42
 Carter, S., 124

 Castro, M. M., 124
 Caveness, W. F., 89-91, 89t, 90ts, 91t, 98-99, 99t, 103
 Chao, D. H.-C., 29-30, 32-37, 48-50, 76
 Chaudhry, M. D., 147, 154
 Christian, —, 65
 Citronblatt, —, 159, 170t
 Clarke, Sir C., 75
 Clemmensen, C., 24-25, 24t
 Clusz, K., 109
 Cobb, W., 124
 Collins, A. L., 134-135, 140, 282
 Conn, J. C. M., 142-143
 Cooper, J. E., 44
 Cosgrove, J. B. R., 9-10
 Cotte, Mme., 40
 Counts, S., 116-117, 117t
 Courjon, J., 40, 96, 100-102
 Craig, W. S., 30-32
 Credner, L., 89-91, 89t, 90ts, 91t, 93t, 94-96, 99t, 104
 Currier, R. D., 65-66, 70-71, 70t, 71t

 Danielsen, J., 83
 Dawson, S., 142-143
 Dekaban, A., 147
 Delaveleye, R., 137-138
 Dennerll, R. D., 135-136, 153
 Denny-Brown, D., 89-92, 89t, 90t, 91t, 96
 Dixon, W. J., 237
 Dominian, J., 10, 17
 Doose, H., 49, 53, 64-65, 70t, 81-82
 Dummermuth, G., 78-79, 85
 Dusaucy-Bauloye, A., 77, 83-84

- Ekholm, E., 45-47, 45t, 47t
 Engel, G. L., 151
 Evans, J. H., 89-91, 89t, 90t, 91ts, 93-94, 93t, 96, 98-100, 99t
 Fabing, H. D., 3
 Falconer, M. A., 117-118, 117t, 118t, 120-121
 Farmer, T. W., 169
 Faxén, N., 24-25, 24t, 32, 47t
 Fegersten, L., 117-118, 117t, 118t, 121
 Fenyes, I., 117-118, 117t, 118t, 120-121
 Fetterman, J. L., 143
 Fischgold, H., 40-41
 Flanigin, H., 117t
 Folsom, A., 136-137
 Fonner, D., 141-142
 Ford, F. R., 33
 Forster, F. M., 3
 Fowler, M., 45-46
 Fox, J. T., 140-141
 Frantzen, E., 10-11, 11t, 17, 45, 45t, 47t, 52-53
 French, L. A., 124
 Friderichsen, C., 44, 46-50, 47t
 Friedmann, M., 54-55
 Fukuyama, Y., 24-26, 24t, 29, 31-33, 40, 46, 48-49, 78
 Guhagan, L., 143-145
 Gastaut, H., 33-34, 84
 Gibberd, F. B., 69-71, 71t
 Gibbs, E. L., 37-40, 73-74, 151-154
 Gibbs, F. A., 38-40, 73-74, 117t, 120, 122-123, 151-154
 Gliddon, W. O., 89-91, 89t, 90t, 91t, 93-94, 93t
 Glithero, E., 148-150
 Goldensohn, E., 124
 Goodall, R. J., 124
 Gowers, W. R., viii, 8, 10-14, 34, 156-157, 164, 219-220, 219t
 Green, J. R., 117t, 119, 121
 Greenstein, L., 17
 Grinker, R. R., 9-11
 Grosz, W., 4t, 9, 11, 32, 160
 Gruetznar, A., 85
 Gruble, H. W., 159, 170t
 Guillaume, J., 117t
 Gunná, J., 117t, 120
 Guttmann, I., 8-9, 159
 Habermaas, —, 4t, 9-10, 14, 170t
 Hahn, —, 159
 Halstead, H., 137
 Harris, R., 41-42, 82-83, 85
 Hedenstroem, I., 9, 11-15, 17-18
 Henderson, P., 136
 Henriksen, P. B., 169-170, 170t, 329
 Herlitz, G., 45, 45t, 47, 47t
 Hertoft, P., 66-67, 70t
 Hess, R., 24-25, 24t, 27, 31-33, 39-40
 Hill, D., 121, 153
 Himler, L. E., 9, 14
 Hippocrates, 9, 15, 51, 106, 220
 Hoefer, P. F. A., 15
 Holowach, J., 62-63, 70t, 71t
 Horsley, Sir V., 106-108, 110-111t, 116
 Horstmann, W., 47-49, 47t
 Hrbek, A., 48-49
 Hunter, R. A., 165
 Hyllested, K., 9
 Illingworth, R. S., 147
 Isler, W., 39-40
 Jabbour, J. T., 24, 24t, 30, 32-33, 40
 Jablon, S., 89-93, 89t, 90ts, 91t, 93t, 96-101, 99t, 103, 139-140, 139t
 Jackson, J. H., 343
 Janz, D., 13, 60, 70t, 71t, 133-134
 Jasper, H. H., 113-114, 114t, 116, 118t, 120-121
 Jeavons, P. M., 75, 77, 79-80, 84-85
 Jenkins, C. D., 152
 Jennett, W. B., 89t, 91t, 95-96
 Jennings, C. C., 48-50
 Joedicke, P., 159, 170t
 Juul-Jensen, P., 4t, 10-15, 11t, 17-18, 20-21, 132
 Keith, H. M., 24-25, 24t, 31-33, 138, 168
 Kennedy, W. A., 121
 Kiørboe, E., 4t, 9-11, 11t, 13, 15, 17, 18-19, 24t

- Kirstein, L., 4t, 8, 13
 Koch, Fr., 85
 Krohn, W., 166-170, 170t
 Kugelmass, I. N., 145
 Kuhlo, W., 60, 68-71, 70t, 71t
- Lamontagne, A., 40-41
 Landolt, H., 17-18, 151
 Larramendi, L. M. H., 124
 Laufer, M. W., 10, 15-16
 Lees, F., 63-64, 70t
 Lempp, R., 9, 12
 Lennox, M. A., 46, 48-50, 220
 Lennox, W. C., 11, 14, 24t, 27-28, 44, 48-49, 54, 61-62, 71t, 73, 76, 126, 133-134, 140, 153, 162-163, 168, 205-206
 Levy, D. M., 44, 48-49
 Lewin, W., 91t
 Liss, H. R., 89-91, 89t, 90ts, 91t
 Liversedge, L. A., 63-64, 70t
 Livingston, S., 23, 33-34, 45t, 47-53, 47t, 67-68, 70-71, 71t, 76-77, 138, 150, 156
 Locock, C., 75
 Lorgé, M., 4t, 11-12, 14, 131-132
 Low, N. L., 80
 Lund, M., 208, 211, 346
 Lundervold, A., 24-25, 24t, 30, 32-33, 40
- MacKinnon, J. M., 30-32
 Maki, Y., 96
 Marshall, C., 101
 Masquin, A., 96, 100-102
 Matera, R., 124
 Matthes, A., 80, 133
 Matthiolius, ---, 108-109, 112-113
 McGreal, D. A., 48-50
 McKissock, W., 124
 Melchior, J., 44, 46-50, 47t
 Melin, K-A., 45t, 48
 Merritt, H. H., 9-12, 11t
 Messina, C., 17, 19
 Meyers, R., 115-116, 123
 Michaux, M. L., 147, 154
 Miller, J. G., 176
 Millichap, J. G., 45t, 47t, 51-53
- Moltke, M. L., 24-25, 24t
 Monrad, S., 23-25, 24t, 30, 32
 Morris, A. A., 117t
 Munson, J. F., 157-158, 160, 167, 170t
 Muskens, L. J. J., 10-12, 14, 32, 130-131, 159, 170t
- Niemineva, K., 45-47, 45t, 47t
 Northfield, D. W. C., 117-118, 117t, 118t
- Obrador, S., 124
 Oppenheim, H., 10, 14
 Ostmann, ---, 160
 Otto, O., 163-165
- Paal, G., 62, 65, 68, 70-71, 70t, 71t
 Pache, H-D., 48, 80, 85
 Paillas, J. E., 117-118, 117t 118t, 120-121
 Paine, K., 117t
 Pakkenberg, H., 9
 Pampiglione, G., 124
 Paskind, H. A., 148
 Passouant, P., 42
 Patrick, H. T., 44, 48-49
 Patterson, H. A., 141-142
 Pauli, L. L., 78, 85
 Peiffer, J., 167-168, 170t
 Penfield, W., 113-114, 114t, 116-117, 117t, 120
 Perkins, R. F., 10, 15-16
 Peterman, M. G., 45, 45t, 48-50
 Petersen, D., 33
 Petit-Dutaillis, D., 117t
 Phillips, G., 89-91, 89t, 90t, 91t, 94, 98, 102
 Picaza, J. A., 117t, 120
 Pigott, A. W., 162
 Plum, P., 78
 Polilisch, K., 54, 58, 70t
 Pond, D. A., 147, 154, 165-166
 Prichard, J. S., 41-42, 48-50
 Probst, C., 4t, 10-11, 13, 17, 150-151
 Pruyser, P. W., 136-137
 Pussep, L., 112-113, 115-116, 123
- Quadfasel, A. F., See "Folsom."

- Ragonese, G., 17, 19
 Ranheim, B., 11
 Ransohoff, J., 124
 Raphael, T., 9, 14
 Rasmussen, T., 116, 118-120, 117t, 118t, 122, 124
 Rawling, L. B., 89-91, 89t, 90t, 91t
 Redlich, E., 94, 100
 Romano, J., 151
 Rodin, E. A., 102, 153, 186, 231-232
 Rosenthal, C., 59-60, 70t
 Russell, W. R., 89-90, 89t, 90ts, 93-94, 93t, 97
- Sargent, P., 89t
 Sauer, H., 55-58
 Sauveur, J. P., 137-138
 Scheetz, D. G., 117t, 119, 121
 Schinnerling, W., 47-49, 47t
 Scheffner, D., 64-65, 70t
 Schmidt, G., 81, 85
 Schorsch, G., 9, 11-15, 17-18
 Schroeder, —, 55
 Schwab, R. S., 16
 Schwade, E. D., 163-165
 Selbach, H., 9, 12
 Serafetinides, E. A., 10, 17, 117t, 118t, 120-121, 148
 Shanks, R. A., 51
 Simmel, M. L., 116-117, 117t
 Slater, E., 148-150
 Sorel, L., 77, 83-84
 Spiegel, E. A., 123
 Spratling, —, 157, 170t
 Stamps, F. W., 77-78
 Steelman, H. F., 116-117, 120
 Steinsiek, H.-D., 160-162, 167, 170t
 Stevenson, W. E., 101
 Stolecke, H., 80, 85
 Stoller, A., 151-152
 Stone, T. T., 4, 5
 Strobos, R. R. J., 4t, 8-14, 20
- Sullivan, E. B., 143-145
 Symonds, C. P., 97-99
- Tenny, J. W., 146
 Thom, D. A., 44
 Tibbles, J. A. R., 41-42
 Tilmann, —, 109-112
 Tissot, —, 61
 Tizard, J. P. M., 41-42
 Toman, J. E. P., 16
 Trojaborg, W., 78
 Trolle, E., 3-4, 4t, 8, 10-11, 11t, 13
 Turner, E., 122
 Turner, W. A., 3-4, 4t, 8-14, 126-131, 128t, 147-148, 157, 170t, 339
- Ulrich, J., 19
- Wagstaffe, W. W., 89-91, 89t, 90t, 91t, 97-98, 104
 Wakana, A., 19
 Walker, A. E., 89-93, 89t, 90ts, 91t, 93t, 96-104, 99t, 139-140, 139t
 Watson, C. W., 89-90, 89t, 90t, 93-94, 93t, 99-101, 99t
 West, W. J., 74-75, 302
 Whitty, C. W. M., 89-90, 89t, 90ts, 93-94, 93t, 97
 Wilkins, L., 24t, 26-27, 31
 Williams, D., 102
 Woodcock, S., 9-10
 Woring, E., 117t, 119-120
 World Health Organization, 30
- Yacorzynski, G. K., 145-146
 Yahr, M. D., 5, 11-13, 11t, 20
 Yannet, H., 23, 28-29, 40
- Zellweger, H., 46-49, 47t
 Zenker, C., 42
 Zimmerman, F. T., 135, 135t
 Zimmerman, H. M., 45-46

SUBJECT INDEX

A

- Abiotrophy, 161, 167
- Absence, 196, 211, 299 (see also Petit mal)
- Accident, as cause of death, 158, 162, 166, 168, 170
- Adolescence, improvement during, 35, 39
- Age
 - analysis of variance, findings, 212-216
 - at onset of
 - illness, 8-10, 23-43, 48, 348
 - analysis of variance, findings, 216-218, 301-302
 - EEG abnormality, 16, 17, 19
 - institutionalized patients, 185, 320, 323, 324
 - intelligence, 134-137, 301-302
 - mental deterioration, 128, 129, 148
 - recurrence of seizures, 21
 - seizure control, 8, 9, 23, 27, 29-31, 41
 - infantile spasms, 76
 - petit mal
 - and incidence of grand mal, 61-63, 65-68
 - treatment
 - results, infantile spasms, 80, 82
 - at time of injury
 - and incidence of posttraumatic epilepsy, 92
 - EEG as prognostic indicator, 19
 - intelligence, 134-136, 208, 211, 212t, 216, 228, 299
- Age (continued)
 - status epilepticus, 213t, 216
 - surgical results, 109
- Aggression, 122
- Akinetic seizures (see Seizures, types)
- Alcohol, 92, 100
- Annesia
 - for seizures, remission, 11, 28
 - posttraumatic, and incidence of epilepsy, 91, 91t, 95
- Anthropometric studies, 148
- Anticonvulsant treatment
 - borax, 24
 - bromides, 4, 5, 24, 27, 97, 100
 - Dilantin, 5, 16, 52, 63, 68, 69
 - diones, 63
 - discontinuation of, 18, 20, 21, 38, 42, 182, 197
 - febrile convulsions, prevention of, 50-53
 - formula to predict success of, in the hospital (Lafayette Clinic), 237-238, 240-254
 - intelligence and, 142, 143, 147
 - ketogenic diet, 25, 27, 76-77
 - Mogadon, 86
 - pharmaceutical industry and, 233
 - phenobarbital, 4, 5, 24, 27, 50, 51, 58, 63, 68, 69, 97, 100
 - response to, 16, 208, 348
 - age, 212
 - employment, 315
 - Performance IQ, 298
 - steroids, 77-86
 - succinimides, 63
 - Tridione, 16

- Anticonvulsant treatment (continued)
 Valium, 86
- Asphyxia (see Death, cause of)
- Atrophy, cerebral, 321t, 322
 bilateral, results of surgical treatment, 121
 recurrence of seizures, 21
 seizure control, 15
- Aura
 in petit mal, 62
- B**
- Barbiturate fast activity, EEG
 results of surgical treatment, 121
- Behavior, 263-276
 analysis of variance, findings, 271-274
 deterioration of, 264, 267
 disturbances, 224, 263
 age, 216
 brain damage, 33, 191, 265, 266, 268, 274, 276
 EEG, 151, 265, 271
 employment, 315, 316
 febrile convulsions, 46
 improvement of, 263, 267
 incidence of, 263, 267
 intelligence, 147, 265, 267, 269, 270, 274, 276
 memory, 269
 MEC-A formula for predicting, 274-276
 neurological examination, 268, 274
 organic mental changes, 269
 psychiatric illness, diagnosis of, 268
 psychological tests
 organic findings on, 267, 268, 269, 274
 psychotic tendencies on, 265
 school performance, 265, 268, 274
 seizure(s), 264, 265, 267, 271
 occurring around menstrual period, 272
 psychomotor, 206, 270
 remission of, 264, 266, 273, 274
 state, 264
 social factors in, 266, 268
 sociopathic or antisocial, 270
- Bender-Gestalt Test, 203 (see also Psychological tests)
 behavior disturbances, 268
 remission of seizures, 203
- Benton Visual Retention Test (see also Psychological tests) and EEG, 152
- Birth
 difficulty during delivery
 febrile convulsions, 49, 53
 incidence of posttraumatic epilepsy, 92
 injury, 29, 31, 32, 49, 53, 211, 322
 order
 incidence of posttraumatic epilepsy, 92
 presentation, 32
 weight, 32, 41
- Brain
 damage, 68, 73, 322
 age at onset of
 deterioration of intelligence, 147
 degree of, and posttraumatic epilepsy, 89
 diffuse
 surgical treatment results, 121
 epilepsy associated with, 203, 204, 347
 febrile convulsions leading to, 45
 learning difficulties, school grades, 279, 281
 site and extent of
 deterioration of intelligence, 147
 treatment results, infantile spasms, 82
 overt local changes in
 surgical treatment results, 109, 121
 tumor, 9
 as cause of death, 162, 166, 168, 170, 193
- Bromides (see Anticonvulsant treatment)
- C**
- Caro State Hospital for Epileptics, 181, 192, 193, 224, 320-325 (see also Institutionalized patients)
- Centrencephalic seizures (see Seizures, types)

Cerebral
 defect, congenital, 29, 32, 323
 infection, 32, 211, 322, 324
 Cerebrovascular disease, 14
 Combination of seizures (see Seizures)
 Concentration, 134
 Confusional episodes, 154
 Cyanosis
 during seizures, 30

D

Death (see also Mortality and Life expectancy)
 age at, 157, 159, 161, 162, 167, 168, 170, 193, 326, 327t, 327
 cause of, 194t
 accident, 158, 162, 166, 168, 170
 asphyxia, 156, 161
 brain tumor, 162, 166, 168, 170, 193
 drowning, 156, 162, 164, 166, 167, 168, 327
 encephalitis, 166, 168
 pneumonia, 158, 161, 166, 167, 168
 pulmonary disorders, 158, 159, 161, 162
 seizure(s), 156, 158, 159, 160, 161, 162, 165, 168, 194t
 status epilepticus, 157, 158, 159, 160, 161, 162, 165, 166, 167, 168, 169, 170, 194t
 suicide, 160, 163, 166, 170
 certificates, 163, 164, 165, 170, 327-329
 childhood epilepsy, 168
 duration of illness, 160, 167, 168, 193
 EEG, 170
 etiology of epilepsy, 162, 167, 169
 seizure(s)
 frequency of, 168, 169, 170
 type of, 169
 sex, 169
 sudden, 156, 166, 167
 Dementia, 15, 84, 126, 128, 130, 131, 132
 posttraumatic epilepsy, 98
 related to seizure frequency, 129
 temporary, 147, 307-313

Depression, 270
 Diagnosis, differential
 of petit mal epilepsy, 55, 58, 70
 of posttraumatic epilepsy
 EEG family studies, 87
 Dilantin, 5, 16, 307
 electroencephalogram, 16
 febrile convulsions, prevention of, 52
 petit mal epilepsy, 63, 68
 Disability
 posttraumatic epilepsy, 98, 99
 Driver's license, 261
 Drowning (see Death, cause of)

E

Electrical stimulation, 107, 108, 113
 Electrocorticography, 113, 122
 Electroencephalogram, 15-19, 34-42, 183, 186, 198-200, 237, 245, 322
 age at onset of illness, 16, 17, 19, 41, 190t, 191, 218
 alpha rhythm, 151, 271, 322
 an independent factor, 188-189, 190, 190t
 anticonvulsant treatment
 Dilantin, 16
 discontinuation of, 20, 42
 response to, 16
 Tridione, 16
 as prognostic indicator,
 posttraumatic epilepsy, 101, 102
 background rhythms, 15, 17, 18, 19, 35, 36, 37, 134, 151, 152, 153, 184, 188, 190, 203, 205, 231, 234, 265, 283, 322
 petit mal, 66, 69
 behavior disturbances, 265, 271
 Benton Visual Retention Test, 152
 bilateral disturbance, 41, 120, 228
 criteria related to surgical treatment, 121
 discrepancy between clinical findings and EEG, 15, 16, 17, 19, 34
 duration of illness, 17
 dysrhythmia, 19
 monorhythmic, 36
 nonspecific, 17
 encephalopathy, 16
 family history of epilepsy, 16, 17

Electroencephalogram (continued)

- family studies in
 - differential diagnosis
 - posttraumatic epilepsy, 87
 - febrile convulsions, 48, 49, 51
 - focal, 16, 18, 19, 20, 36, 37, 39, 184, 199, 231
 - abnormalities in patients with
 - petit mal, 69
 - slow spike and waves
 - posttraumatic epilepsy, 102
- focus(i)
 - epileptogenic, 115, 123, 343-344
 - laterality of
 - Wechsler Test, 137, 153
 - left sided, 137, 323
 - migration of, 37
 - multiple, 123
 - occipital, 37, 39
 - spike or sharp wave, 37, 40, 41, 113
 - temporal, 132, 153
 - anterior, 121
 - remission, 37, 39
 - independent bitemporal, 19
 - left, 137, 232
 - age related, 216
 - mid-, 36, 37, 38, 39, 121
 - remission, 37, 39
 - right, 137, 216, 232
 - temporal occipital, 37
 - follow-up studies, 15, 16, 17, 18, 19, 37, 38, 39-40, 41, 79, 183-185, 183t, 184t, 197-220, 198t, 199t
 - fourteen and six cycles per second
 - positive spikes, 35, 36, 37, 38, 39, 248
 - head injury, 16
 - heredity of, 345
 - hyperventilation, 15, 250
 - hypsarhythmia, 35, 39, 73, 74, 76, 79
 - improvement of, 16, 17, 18
 - infancy, spells in, 16
 - infantile spasms, 74, 76, 78, 79, 81, 82, 83, 86
 - initial versus follow-up, 17, 18, 34, 183-185, 197-200
 - intelligence, 41, 42, 147
 - interictal, 19

Electroencephalogram (continued)

- larval attacks
 - posttraumatic epilepsy, 102
- learning difficulties, 279, 283
- mean age at death, 170
- medication withdrawal, 18, 20
- mental and motor development, in
 - prediction of, 35, 41
- mental state, 132, 134, 150-154, 310
- mitten patterns, 153
- multiple spikes followed by wave
 - myoclonic jerks, 73
- normal, 15, 16, 17, 18, 19, 35, 37, 38, 40, 48
 - posttraumatic epilepsy 101, 102
 - review of concept, 151, 293
- normalization of, 38, 39, 40, 42, 51, 77, 83, 101, 102
 - forced, 151
- paroxysmal activity, 15, 17, 18, 20, 36, 37, 206, 231, 234
- Performance IQ, 288
- petit mal, 62, 66, 197 (see also *Petit mal*)
 - status, 307-313
 - variant, 34, 39, 73, 134, 152
- photic driving response, 231
- IQ, 288t
 - learning difficulties, 283
- post injury
 - posttraumatic epilepsy, 92
- posterior emphasis of spike wave
 - discharges
 - petit mal, 66
- postoperative, as prognostic indicator, 120-121
- preoperative, as prognostic indicator, 120
- seizure(s)
 - akinetie, 73
 - centrencephalic, 17
 - combination of, 16
 - focal, 17, 19
 - frequency of, 16, 17
 - improvement of, 17
 - neonatal, 41
 - patterns, 183t, 184, 198t, 199, 200, 201t, 202, 203, 204t, 208, 231

- Electroencephalogram (continued)
- seizure(s) (continued)
 - remission of, 16, 37-42, 188, 201, 228-231
 - subclinical
 - posttraumatic epilepsy, 102
 - threshold, 232-233, 345
 - time of day, 17
 - serial recordings
 - posttraumatic epilepsy, 102
 - single versus multiple spike discharges
 - petit mal, 66
 - sleep recording, 38, 41, 231
 - sleep-waking cycle, 13
 - slow spike and wave, 35, 73
 - spike wave activity, 40, 62, 285
 - Full Scale IQ, 285
 - learning difficulties, 283
 - Performance IQ, 154, 286, 287t, 288
 - three cycles per second, 34, 35, 36, 63, 153
 - surgical treatment, 120, 121
 - two to three cycles per second spike
 - wave bursts, 66, 73, 152
 - work ability, 16, 315t, 316t
 - Employment, 122, 124, 153, 203, 205t, 314-319
 - anticonvulsant treatment
 - patient's response to, 315
 - behavioral difficulties, 315, 316
 - intelligence, 315, 316, 318, 319
 - organic mental changes, 315, 316, 318, 319
 - personality difficulties, 316, 318, 319
 - seizure(s)
 - frequency, 315, 318
 - remission, 318
 - series of, 315
 - state at follow-up, 317
 - types, 315
 - state
 - analysis of variance, findings, 315-317
- Encephalitis, 28
- as cause of death, 166, 168
- Encephalopathy
- diffuse, 30, 36
- Encephalopathy (continued)
- EEG abnormality, 16, 36
 - infantile, 14
- Environment, social
- deterioration of Full Scale IQ, 294
- Environmental strain
- remission, 27
- Epilepsia Nutans, 74
- Epilepsy
- associated with brain damage, 203-204, 204t, 346
 - childhood, 23-85, 180-192, 199, 200, 204, 211-218, 263-267, 277-280, 281, 306, 347, 348
 - mortality, 160, 162, 168, 169
 - constitutional factor in, 345
 - definition of, 100, 181
 - etiology of, 30, 93, 103, 196, 206, 211, 231, 234, 323, 344, 345
 - analysis of variance, findings, 323
 - birth injury, 29, 31, 32, 196, 211, 322, 324
 - cerebral
 - infection, 32, 196, 211, 322, 324
 - malformations, 29
 - family history, 14, 17, 29, 30, 42, 48t, 126, 128, 132, 196, 211, 218, 232, 344
 - EEG, 16, 17, 345
 - abnormality, 16
 - febrile convulsions
 - recurrence rates, 45
 - intelligence
 - deterioration of, 128, 132, 147, 300
 - petit mal
 - changing to grand mal, 62, 66, 67
 - course of illness, 62, 65, 66, 67, 69
 - incidence of, 62, 66
 - posttraumatic epilepsy
 - incidence of, 92, 94
 - recurrence of seizures, 20, 21
 - remission of seizures, 14, 27, 29, 211, 232, 234
 - febrile convulsions as first symptom
 - of, 47t, 46-50, 52, 53, 68
 - genetic, 29, 30

Epilepsy (continued)

- grand mal (see Grand mal)
 - idiopathic, 29, 30, 99, 101, 283, 344
 - immediate, 95
 - "malignant," 324
 - pathophysiology of, 220, 342-344
 - petit mal (see Petit mal)
 - posttraumatic (see Posttraumatic epilepsy)
 - prevention of, 51, 52, 53
 - psychomotor-temporal lobe (see Seizures, types)
 - symptomatic versus cryptogenic disorder, 13, 30, 31, 32, 79, 80, 87, 96, 344
 - traumatic, Jacksonian, 109
- Epileptic Cry
- seizure control, 186t, 188
- Epileptic equivalents, 12, 42, 115, 138
- Epileptic facies, 126, 127, 130
- Epileptic personality change, 58
- Epileptische Wesensveraenderung, 132-133, 151

F

Facies Epileptica, 126, 127, 130

Family history

- of epilepsy (see Epilepsy, family history)
- of mental retardation, 279
- of nervous disorders
 - incidence of posttraumatic epilepsy, 92

Febrile convulsions, 44-53, 343

- as first symptom of epilepsy, 46, 47, 47t, 51, 53
- behavior disorders in, 46, 274
- benign versus atypical, 47-50, 53
- follow-up studies, 45t, 46, 47, 47t, 51, 52, 53
- incidence of, 44
- leading to brain damage, 45
- mental development in, 46
- prevention of, 50-53
- psycholability in, 46
- recurrence rates, 45, 45t, 51, 52
 - related to family history, 45, 53
- speech disorders in, 46

Feeding, problems in infancy

behavior disturbances, 274

Fevers, excessively high

learning difficulties, 279

Flexion spasms, 74

Focal seizures (see Seizures, types)

Follow-up

- criteria, 115
- surgical treatment results, 116
- duration of
 - remission of seizures, 4, 5, 6t
 - treatment results, infantile spasms, 78-79

G

Grand mal

petit mal changing to

- incidence of, 60, 62, 65, 67, 68, 69, 71t
- mentality, 66
- neurological examination, normal, 65, 66
- prophylactic treatment, 63, 68, 69
- sex, 62, 65, 69
- pyknolepsy, as a prelude to, 60
- seizure(s), 11, 11t, 12, 17, 28, 29, 30, 32, 65, 66, 67, 68, 69, 122, 162, 169, 195t, 322t
- intelligence and, 134, 135, 135t, 136, 137
- remission of, 11, 11t, 12, 28, 29, 30, 32
- variant
 - focal, induced by Megninide, 232
 - non-focal, 231

H

Handedness

remission, posttraumatic epilepsy, 103

Head injury (see Posttraumatic epilepsy)

Hemiplegia, acute infantile, 33

Heredity (see Epilepsy, etiology, family history)

H.H.E. Syndrome, 33-34

Hormonal treatment, 82 (see also Steroids)

Hypsarrhythmia, 35, 39, 73, 76-86, 185, 347 (see also Infantile spasms)

I

Illness

- age at onset of, 216-218, 320, 321t, 324, 325t, 348
- remission of seizures, 27, 29, 31, 41, 217, 218
- duration of, 218, 320, 346
 - age at death, 160, 167, 168, 193, 194t
 - electroencephalogram, 17
 - Full Scale IQ, 285
 - intelligence, 134, 137
 - learning difficulties, 279
 - mental deterioration, 128, 132
 - Performance IQ, 286
 - personality changes, 132
 - prior to treatment
 - results, infantile spasms, 81-86
 - remission, seizures, 10, 27, 29, 135t, 186t, 188, 201t, 203, 207, 208, 218, 219t, 223, 223t, 234
 - severity of, 228, 320, 323, 324, 346
 - remission of seizures, 11, 27, 234

Inattention, 147

Infancy

- feeding problems in
 - behavior disturbances, 274
- isolated convulsions in, 280
 - behavior disturbances, 266-267, 274
 - learning difficulties, school grades, 280, 282-283
- massive myoclonic jerks of, 74
- onset of seizures in
 - behavior disturbances, 265
 - Full Scale IQ, 285
 - learning difficulties, school grades, 282-283
 - Performance IQ, 286
- spells in,
 - EEG abnormality, 16

Infantile encephalopathy, 14

Infantile spasms, 73-86, 185

- age at onset, 76
- a medical emergency, 86, 347, 348
- combined with other seizures, 76, 79
- course of illness, 76, 77, 79, 80, 81, 83

Infantile spasms (continued)

- electroencephalogram, 74, 76, 77, 79, 81, 82, 83, 85, 86
- etiology, 76, 79
- follow-up studies, 78-83
- incidence of, 76, 78
- intelligence, 76, 81, 147
- mental
 - deterioration, 76, 77, 82
 - development, 76, 80, 81, 82, 83, 84, 85
 - improvement, 77-80
 - retardation, 76-80, 83, 85
 - prevention of, 77, 82, 85, 347, 348
- mortality, 77, 79, 80, 81
- relapse after treatment, 78, 79, 80, 83, 85
- series of seizures, 76
- treatment
 - ACTH, 77, 78, 80, 81, 82, 83, 84
 - Cortisone, 78, 80, 81
 - ketogenic diet, 76
 - Mogadon, 86
 - Prednisolone, 79, 80
 - Prednisone, 81
 - Valium, 86
- treatment results, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85
 - age at onset of treatment and, 80, 82
 - brain damage (pre-existing) and, 82
 - duration of illness prior to treatment, 81, 82, 83-86
 - length of follow-up, 78

Injury

- age at time of
 - incidence of posttraumatic epilepsy, 92
- civilian versus war
 - incidence of posttraumatic epilepsy, 96
- during seizures, 208, 228, 237, 320
- missile versus non-missile, 103
- pre and/or perinatal
 - etiology of epilepsy, 322-324
- severity of posttraumatic epilepsy, 103

- Insanity, 127-128
- Institutionalized Patients, 12, 14, 15, 33, 157, 158, 159, 160, 165, 166, 184, 185, 190, 190t, 191, 194, 224, 233, 320-325, 321t (see also Caro State Hospital for Epileptics)
 - age at onset of illness, 185, 320
 - analysis of variance, findings, 321-323
 - intelligence, 126, 127, 131, 140, 141, 145, 322t, 324
- Intellectual
 - deficit
 - discontinuation of anticonvulsant treatment, 42
 - deterioration
 - MEC-A formula for predicting, 303-306
- Intelligence, 126-155, 203, 228, 277-313, (see also Dementia, IQ, Mental deficiency, Mental deterioration, Mental growth, Mental retardation, and Mental state)
 - age, 134, 135, 213, 213t, 216, 299
 - at onset of illness, 134, 135, 137, 138, 301, 301t, 302
 - anticonvulsant medication, 142, 143, 147, 298
 - behavior disturbances, 265, 267, 276
 - deterioration of, 142, 143, 147, 347
 - discriminant function analysis, 302-306
 - seizure state, 291
 - duration of illness, 134, 137, 285
 - EEG abnormalities, 134, 147, 285
 - employment, 315-316
 - improvement of, 141-144, 147
 - infantile spasms, 76, 79, 85
 - inpatients versus outpatients, 131
 - learning difficulties, school grades, 281
 - of healthy siblings, 140, 142
 - private versus clinic patients, 134
 - posttraumatic epilepsy, 135, 139, 139t
 - remission, seizures, 14, 27, 32, 203, 228, 294-298
 - posttraumatic epilepsy, 101, 139
- Intelligence (continued)
 - seizure(s)
 - combination of, 133, 137, 283, 292t, 303t
 - frequency, 133, 136, 137, 138, 141, 142, 143, 146, 147
 - severity of major, 133, 137, 142, 143, 146
 - symptomatic versus genetic disorder, 134, 135, 135t, 137, 138, 145
 - type(s)
 - grand mal, 134, 135-137
 - petit mal, 29, 67, 133, 134, 135, 137, 138, 140
 - psychomotor-temporal lobe, 134, 136
 - serial tests, 140-146
 - variability of, 141, 142, 143, 144
- IQ, Full Scale
 - age at onset of illness, 285, 301, 302
 - behavior disturbances, 269t
 - changes in
 - analysis of variance, 290-294
 - comparison of initial to follow-up, 289-298
 - and seizure frequency, 294-298
 - EEG spike wave activity, 285
 - etiology, 291
 - fluctuations in, 293
 - neurological examination, 284
 - school performance, 283
 - seizure(s)
 - duration of disorder, 285
 - during first year of life, 285
 - focal minor motor, 285
 - frequency, 285, 292t
 - state at follow-up, 285, 294-296
 - social environment, 294
- IQ Performance, 134, 135, 135t, 136, 152, 290, 295-300
 - behavior disturbances, 269, 274
 - changes in
 - analysis of variance, 298-300
 - comparison of initial to follow-up, 295t
 - and seizure frequency, 295-297
 - EEG, 152, 286, 288
 - neurological examination, 286

IQ (continued)

- organic findings on psychological tests, 286
- seizure(s), 299, 299t
 - during infancy, 286
 - focal minor motor, 286
 - frequency of, 298
- sex, 299
- IQ, Wechsler, Subsections, 289t
 - arithmetic, 134, 290
 - block design, 134, 152, 153
 - comprehension, 134, 136, 153
 - digit span, 134, 136, 153
 - digit symbol, 136, 152, 290
 - object assembly, 152, 153, 290, 302
 - picture arrangement, 290
 - picture completion, 153, 290
 - Porteus Maze, 152
 - similarities, 153
- IQ, Verbal, 134, 136, 152, 298, 301
 - age at onset of illness, 301
 - comparison of initial to follow-up, 289-290
 - seizure frequency, 294-298

K

- Ketogenic diet, 25, 27, 34, 76-77

L

Lafayette Clinic

- formula to predict success of anti-convulsant treatment in hospital, 237-239, 240-254, 261
- inpatient treatment results, 225, 226t, 227t, 228, 229t, 230t, 231, 231t
- outpatient treatment results, 220-223
- Learning difficulties, 282t
 - brain damage, 279, 281
 - electroencephalogram
 - background rhythms, 279, 283
 - photie driving response, 283
 - spike wave activity, 283
 - fevers, excessively high, 279
 - illness, duration of, 279
 - incidence of, 277, 280
 - infancy, isolated convulsions in, 280
 - intelligence, 282
 - IQ, Verbal, 282
 - mental retardation, family history of, 279

Learning difficulties (continued)

- neurological examination, 278t, 279
- seizure(s)
 - combination of, 283
 - cyanosis during, 279
 - during first year of life, 281t, 282
 - focal motor, 279
 - psychomotor-temporal lobe, 279
 - state, 277
 - types, 278
- Librium, 307
- Life
 - expectancy, 156, 157, 159, 160, 162, 167, 168, 169, 170, 326-329
 - variability of, 162, 171
 - insurance, 170, 171
- Limbic System, 270

M

- Manual dexterity, 137
- Megimide, 232, 239
- Memory, 126, 127, 129, 134, 136, 137, 141, 147, 149
 - behavior disturbances, 269
- Meningoencephalitis, 14, 32
- Menstrual period, 13, 253
 - seizures occurring around
 - behavior disturbances, 272
- Mental changes
 - age at death, 162, 169
 - behavior disturbances, 269
 - employment, 314-316
 - remission of seizures, 6, 8, 202
- Mental deterioration, 58, 126, 128t, 131, 136, 144, 145, 147, 152
 - age at onset of seizures, 128-129, 147-148
 - constitution, 148
 - duration of illness, 128
 - heredity, 128, 132, 147
 - posttraumatic epilepsy, 98, 101
 - seizure(s)
 - arrest of, 130, 132
 - combination of, 129, 133, 134
 - frequency of, 129, 132, 133, 147
 - occurring in series, 130
 - sex, 127, 128t
 - stigmata of degeneration, 130

- Mental development, 29, 30
 EEG in prediction of, 35, 36, 41, 42
 febrile convulsions, 46
 infantile spasms, 75, 76, 80-85
 petit mal, 133
- Mental growth, 145, 306, 307
- Mental retardation, 29, 31, 33, 36, 46
 definition of, 138
 family history of
 learning difficulties, school grades, 279
 incidence of, 133, 138
 infantile spasms, 75, 76, 80, 83
 petit mal, 62, 68
 prevention of
 infantile spasms, 77, 82, 85, 347, 348
 seizure frequency, 133
- Mentality, 14, 30, 126, 132
 EEG, 132, 134, 150-154
 improvement of, after surgical treatment, 107
 petit mal, 55, 59, 63, 66
 changing to grand mal, 66
 remission, seizures, 6, 8, 23
- Mesantoin, 309
- Metrazol, 233
- Michigan Department of Health, 326-329
- Michigan Epilepsy Center
 data coding, 180, 186-188
 description of services, 179, 180
 first follow-up study, 180-192, 263-267, 277-280, 320
 formula for predicting
 behavioral disturbances, 274-276
 intellectual deterioration, 303-306
 likelihood of long-term remission, 254-257
 second follow-up study, 192-220, 267-276, 280-300, 314-317, 320
 Vocational Rehabilitation Administration Project, 258-261, 317-318
- Microcephaly, 81
- Mitten patterns in EEG
 mental state, 153
- Mogadon, 86
- Mortality, 156-171, 192 (see also Death, Life expectancy)
- Mortality (continued)
 childhood epilepsy, 160, 168, 169
 infantile spasms, 77, 79, 80, 81
 neonatal seizures, 30, 31, 41, 168, 169
 posttraumatic epilepsy, 104
 rate, 163, 165, 169, 192, 327t
 surgical treatment, 108, 109, 113, 115, 119
- Multiple Sclerosis, 14
- Myoclonic
 jerks, 73, 76
 electroencephalogram in, 73
 of infancy, massive, 74
 seizures, infantile (see Infantile spasms)
- Myoclonus, course of illness
 petit mal epilepsy, 69
- N**
- Nervous disorders, family history
 incidence of posttraumatic epilepsy, 92
- Neurological examination
 behavior disturbances, 269, 274
 Full Scale IQ, 285
 learning difficulties
 school grades, 278, 279
 normal
 petit mal changing to grand mal, 65, 66
- Performance IQ, 285, 286
 seizure control, 14, 27, 191, 203, 211, 228
- O**
- Organic findings
 on psychological tests (see Psychological tests)
- Organic mental syndrome, 202t, 202, 205, 216, 269
- Oxycephaly, 14
- P**
- Paranoid ideation, 122, 149
- Personality
 change, 32, 42, 63, 132, 148-150, 151
 anticonvulsant treatment
 discontinuation of, 42

- Personality (continued)
 change (continued)
 illness
 duration of, 132
 seizure frequency, 132
 difficulties
 employment, 316
 disorder, 270, 293
 seizure control, 27, 32
 surgical treatment results, 121
 Petit mal, 29, 34, 39, 54-72, 76, 101, 113, 197, 205 (see also Pyknolepsy)
 aura, 62
 brain damage, 62, 68
 change to grand mal
 incidence of, 60, 62, 65, 67, 68, 69, 71t
 age at onset of petit mal, 61, 63, 65, 66, 67, 68
 frequency of petit mal, 66
 heredity, 62, 66, 67
 mentality, 66
 neurological examination, 66
 prophylactic treatment, 65, 68, 69
 sex, 62, 65, 69
 time of onset of treatment, 64
 change to psychomotor-temporal lobe seizures, 62, 65
 combination of seizures, 61, 62, 63, 64, 68, 69, 70
 course of illness, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70
 age at onset of petit mal, 61, 62, 63, 66, 67, 68
 aura, 62
 brain damage, 62
 early morning seizures, 62
 etiology, 67, 69
 myoclonus, 69
 sex, 62, 65, 67, 69
 differential diagnosis, 54, 58, 70, 113
 electroencephalogram
 background slowing, 66, 69
 focal abnormalities, 69
 follow-up findings, 62, 65, 66, 67, 68, 69
 in differential diagnosis, 54
 Petit mal (continued)
 electroencephalogram (continued)
 in prognosis for
 course of petit mal, 62, 66, 69
 spike discharges
 single versus multiple, 66
 spike wave
 activity, three cycles per second, 54, 62, 63, 68, 69, 70, 71, 101, 153
 bursts
 atypical, 62, 66, 69
 posterior emphasis of, 66
 two to three cycles per second, 66, 152
 family history of epilepsy, 62, 65, 67, 69
 first description of, 61
 incidence of, 61, 62, 67
 intelligence, 29, 58, 66, 67, 133, 134, 135, 135t, 137, 138
 mentality, 58, 63, 66
 mental retardation, 62, 68
 personality change, epileptic, 58
 posttraumatic epilepsy, 101
 propulsive, 74
 puberty
 cessation at, 58, 63, 64
 remission, 12, 28, 29, 30, 32, 60, 62, 63, 64, 65, 66, 67, 68, 69, 70t
 relationship to onset of treatment, 64, 67
 spontaneous, 60, 63
 sex, 60, 62, 65, 66, 67, 69
 status, 58, 68, 69, 153, 154, 307-313
 variant, 34, 39, 73, 152, 153
 with features of automatisms, 65
 Phenobarbital, 4, 5, 24, 27, 38, 51, 58, 63, 68, 69, 97, 100
 febrile convulsions, prevention of, 50, 51
 Pneumoencephalogram, 322
 discontinuation of anticonvulsant treatment, 21
 seizure control, 15
 surgical treatment results, 121
 Posttraumatic epilepsy, 10, 87-105
 definition of, 87, 100
 dementia, 98

- Posttraumatic epilepsy (continued)
 differential diagnosis, 87
 EEG, family studies in, 87
 disability, 98, 99
 early versus late, 95, 96, 97, 97t, 98
 electroencephalogram as prognostic indicator, 92, 101, 102
 frequency of seizures, 98, 100, 101
 incidence of, 88, 89, 89t, 90t, 93, 95, 96
 age at time of injury, 92
 alcohol consumption, 92
 amnesia, length of posttraumatic, 91, 91t, 95
 birth order, 92
 cerebral damage, degree of, 89, 90
 constipation, 92
 delivery, difficulty during, 92
 dural penetration, 88, 90, 91t
 electroencephalogram, post injury, 92
 family history of
 epilepsy, 92, 94
 nervous disorders, 92
 foreign bodies, intracranial, 92, 94
 head injury, previous, 92
 skull fracture
 depressed, 90, 95
 linear, 90
 systemic illness, previous, 92
 unconsciousness, duration of, 91, 92
 wound
 diameter and depth of, 92
 factors leading to delayed healing, 92
 location, 92, 93t, 94
 injuries, civilian versus war, 96
 intelligence, 135, 135t, 139, 139t
 mental deterioration, 98, 101
 mortality, 104
 remission, 14, 99, 99t, 100, 101, 103
 seizure type, 101
 syndrome
 seizure control, 103
 Prophylactic treatment
 febrile convulsions, 50-53
 petit mal changing to grand mal, 63, 68, 69
 Psychiatric difficulties, 14, 202, 202t, 208
 as precursor of overt neurological disease, 292, 293
 Psychological tests (see also IQ)
 Bender-Gestalt Test, 203
 behavior disturbances, 268
 Full Scale IQ, 284, 291
 Performance IQ, 286
 seizure control, 203
 Benton Visual Retention Test and EEG, 152
 data
 surgical treatment results, 121
 organic findings on
 age, 213, 216
 behavior disturbances, 267, 268, 269, 274
 institutionalized patients, 322
 seizure control, 211
 psychotic tendencies
 incidence of behavior disturbances, 265, 265t
 Wechsler Test findings
 and laterality of EEG focus, 137, 153
 weight discrimination test and EEG, 152
 Psychomotor-temporal lobe (see Seizures, types)
 Psychosis, 149, 151, 270
 Puberty
 cessation of petit mal seizures at, 58, 63, 64, 69
 discontinuation of anticonvulsant treatment, 42
 Pyknolepsy, 54-61 (see also Petit mal)
 as prelude of grand mal epilepsy, 60
 case reports, 55-57
 definition of, 55, 58, 59
 incidence of, 58, 60
 relationship to narcolepsy, 55, 58, 59
 retrospective diagnosis, 61
 seizures related to time of day, 58
 sex, 60

 R
 Recurrence rates
 febrile convulsions, 45, 45t, 52

- Recurrence rates (continued)
 of seizures after discontinuation of
 anticonvulsant treatment, 20-21
 Remission of seizures, 3, 4, 4t, 5, 6, 6t,
 7t, 8, 10, 11, 11t, 12, 13, 14, 15,
 16, 23, 24t, 25, 26, 27, 29, 30, 31,
 32, 37, 38, 39, 41, 43, 182, 193,
 196, 205, 220, 221, 222, 225, 233
 age
 at onset of illness, 8, 9, 10, 27, 30,
 31, 41, 185
 chronological, 27, 208, 209t, 211
 anticonvulsant medication
 immediate response to, 208
 behavior disturbances, 32, 266, 273
 cerebral atrophy, 15
 developmental retardation, 32
 electroencephalogram, 15-19, 208
 amount of overall abnormality,
 208, 237
 background frequencies, 186t, 188,
 229t, 231
 focus(i), 37-40, 202t, 229-231t,
 232
 fourteen and six cycles per second
 positive spikes, 37, 39
 seizure patterns, 201t, 202, 208,
 231
 employment, 317
 environmental strain, 27
 epileptic cry, 188
 etiology, 5, 6, 13, 14, 30, 186t, 188,
 211, 231, 234
 birth
 injury 29, 32
 presentation, 32
 weight, 32, 41
 brain injury, 27, 32
 missile versus non-missile, 103
 severity of, 103
 congenital cerebral defect, 29, 32
 diffuse encephalopathy, 30
 meningoencephalitis, 32
 prenatal central nervous system
 insult, 32
 family history of epilepsy, 14, 27,
 29, 30, 232
 follow-up
 duration of, 4, 5, 6t, 24, 25, 26
 Remission of seizures (continued)
 handedness, 103
 illness
 duration of, 10, 27, 186t, 188,
 201t, 202, 207
 intensity of, 11, 27, 228, 234
 intelligence, 14, 27, 32, 103, 203,
 228
 menstrual cycle, 13
 mentality, 6, 8, 14, 15, 23
 mental retardation, 186t, 188, 191
 neurological examination, 14, 27, 191,
 203, 211, 228
 patient sample, 24, 25, 27, 221-223
 personality, 27, 32
 pneumoencephalography, 15
 posttraumatic (see Posttraumatic epi-
 lepsy)
 psychiatric symptoms, 14
 psychological tests
 organic findings on, 203, 211
 school performance, 211
 seizure(s)
 amnesia for, 28
 combination of, 11, 12, 27, 30, 32,
 33, 208, 236
 cyanosis during, 30
 daytime, 13
 duration of, 32, 41
 frequency of, 11, 27, 28, 32, 103,
 228, 236
 injuries during, 208, 228, 237
 nocturnal, 9, 13, 232
 number of, 10, 27
 series of, 76, 208, 228, 237
 type(s), 27, 41, 96
 absence with myoclonic features,
 232
 akinetie, 12, 27, 232
 centrencephalic, 17, 31
 equivalents, 12
 focal, 17, 28, 31
 grand mal, 11, 11t, 12, 28, 29,
 30, 32
 grand mal variant, non-focal,
 231
 minor, 12, 13
 minor motor, focal, 203, 231
 minor motor, non-focal, 231

- Remission of seizures (continued)
- seizure(s) (continued)
 - type(s) (continued)
 - myoclonic, 12
 - infantile, 12, 33
 - petit mal (see Petit mal epilepsy)
 - psychomotor-temporal lobe, 11, 11t, 12, 14, 28, 33, 202, 205, 208, 232, 236, 236t
 - tonic, 33
 - sex, 7t, 8, 202
 - sleep-waking cycle, 13
 - spontaneous, 27, 36, 60, 63
 - status epilepticus, 11, 28, 191, 202, 211, 228
 - temporal lobectomy, 118, 118t
 - time of
 - debridement after head injury, 103
 - onset after head injury, 95, 96, 97, 97t, 98
 - wound
 - complication, 103
 - depth of, 103
 - location of, 103
- Rolandic lesions
- incidence of posttraumatic epilepsy, 92, 93, 93t
- S**
- Schizophrenia-like psychosis, 149-150
- School performance
- behavior disturbances, 265, 268, 274
 - seizure(s)
 - control, 211, 277, 278
 - types, 278
 - Verbal IQ, 286
- Secousses, 74
- Seizure(s)
- arrest of, and mentality, 131, 132, 294-297
 - as cause of death, 156, 158, 159, 160, 161, 162, 165, 168, 194t
 - childhood, isolated in
 - Full Scale IQ, 286t
 - clusters of, 76, 130, 206t, 208, 210t, 213t, 226t, 227t, 228, 235t, 237, 238t, 240t, 241t, 253, 256t, 260t,
- Seizure(s) (continued)
- clusters of (continued)
 - 299, 299t, 303t, 316, 316t, 321t, 324, 325t
 - combination of, 16, 27, 30, 32, 33, 62-70, 182, 190t, 195, 201t, 202, 208, 218, 228, 251, 252, 320
 - intelligence, 134, 135, 135t, 137
 - learning difficulties, school grade, 283
 - mental deterioration, 126, 133
 - Performance IQ, deterioration, 299
 - remission, seizures, 11, 12, 191, 201t, 202, 208, 236
 - with infantile spasms, 75, 79
 - control of, 3, 5, 10, 24, 25, 26
 - in hospital, 233, 234
 - cyanosis during, 30
 - learning difficulties, 279
 - daytime
 - remission rates, 13
 - duration of, 30, 32, 41, 49, 133, 191
 - during infancy, 280, 282
 - Full Scale IQ, 284t, 285
 - learning difficulties, 280, 282
 - Performance IQ, 286
 - early morning, 13
 - petit mal epilepsy, course of illness, 62
 - frequency of
 - anticonvulsant medication
 - discontinuation of, 20, 42
 - dementia, 132
 - EEG abnormality, 16, 17
 - employment, 315, 316, 317, 318
 - Full Scale IQ
 - comparison of initial to follow-up, 295-298
 - deterioration of, 291
 - intelligence, 136, 137, 138, 141, 142, 143, 146, 147
 - institutionalization, 322
 - mean age at death, 168
 - mental
 - deterioration, 51, 129, 133, 148
 - retardation, 133
 - Performance IQ, 295-298
 - personality changes, 132

Seizure(s) (continued)

- frequency of (continued)
 - posttraumatic, 98, 100, 101
 - recurrence after discontinuation of anticonvulsants, 20
 - remission, 11, 27, 28, 32, 103, 228, 236
- Verbal IQ, 295-298
- improvement of, 5, 10, 197, 198, 198t, 211
- infancy, isolated in
 - behavior disturbances, 266, 267, 274
- injuries during, 320
 - remission, 208, 228, 237
- intensity of disorder
 - remission, 11, 228, 234
- major, severity of
 - and intelligence, 137, 138, 146
- menstrual period, occurring around
 - behavior disturbances, 272
- neonatal, 31, 41
 - EEG, 41
 - mortality, 31, 41
- nocturnal, 9, 16, 232
 - remission, 13
- number of
 - remission, 10, 27
- onset in infancy
 - behavior disturbances, 265
 - learning difficulties, 280, 282
- patterns in EEG, 184, 190t, 202, 204t, 205, 208, 211, 235t, 243-251
- Performance IQ, deterioration, 298, 299
- propensity, 206, 206t, 233, 342, 344, 345
- recurrence after discontinuation of anticonvulsant medication, 18, 20, 21, 42
- remission (see also Remission)
 - behavior disturbances, 266, 273
 - employment, 318
- series of, 76, 130, 237, 316, 324
 - mental deterioration, 130
 - Performance IQ, 299
 - remission, 208, 228, 237
- severity of, 133, 137, 146

Seizure(s) (continued)

- threshold, 206, 232, 233, 342, 344, 345, 348
 - EEG and, 345
- time of day of
 - early morning, petit mal, 58-62
 - EEG abnormality, 17
- type(s), 182t, 195t, 225t, 229-231t
 - absence, 195t, 196, 217t, 218
 - Performance IQ, 299
 - school grades, 278
 - akinetetic, 12, 27, 58, 73, 137, 232
 - astatic, 73, 76
 - Blitz-Nick-Salaam (BNS), 74
 - centrencephalic, 17, 31
 - drop, 74
 - equivalents, 12, 115, 138
 - focal, 17, 19, 28, 31, 40, 48, 185, 195t, 281
 - focal minor motor, 34, 74, 76, 185, 190t, 203, 215t, 231, 285, 286
 - Full Scale IQ, 285
 - Performance IQ, 286
- focal motor
 - learning difficulties, school grades, 279
- grand mal, 17, 65, 66, 67, 68, 69, 122, 162, 169, 195t, 322t
- focal variant induced by Megimide, 232
- intelligence, 134, 135, 135t, 136, 137
 - non-focal variant, 231
 - remission, 11, 11t, 12, 28, 30, 32
- infantile spasms, 33, 50, 73-86, 185, 302, 347
- institutionalized patients, 185, 322t
- lightning, 74
- minor
 - remission, 10, 12
- minor motor
 - non-focal, 231
- myoclonic, 12, 73, 197
 - infantile, 12, 33 (see also Infantile spasms)
 - jerks, 73, 76, 197
- petit mal (see Petit mal)
- posttraumatic (see Posttraumatic epilepsy)

- Seizure(s) (continued)
 type(s) (continued)
 psychomotor-temporal lobe, 28, 33,
 34, 39, 120, 122, 151, 153,
 201t, 202, 205, 205t, 208,
 218, 224, 279
 behavior disturbances, 206, 270,
 271
 intelligence, 134, 136
 petit mal changing to, 62, 65, 66
 posttraumatic epilepsy, 101
 remission, 11, 11t, 12, 14, 28,
 33, 39, 202, 205, 208, 232,
 236, 270
 school grades, 278, 279
 treatment, surgical, 116, 122
 salaam, 74, 75
 tonic, 33
 unifocal versus multifocal, 30
- Sex, 130, 141, 202
 febrile convulsions, 49
 mental deterioration, 126, 128t
 mortality, 169
 petit mal, 62, 65, 67
 change to grand mal, 62, 65, 69
 incidence of, 66
- Performance IQ, deterioration of,
 299
- pyknolepsy, 60
 remission, seizures, 6, 8, 201t, 202
- Sleep recording
 EEG, 38
- Sleep-Waking cycle, 13
- Social factors
 behavior disturbances, 266
 environment and Full Scale IQ, 294
- Speech disorders, 32, 46
- Spike wave status, 154, 307-309
- Statistical procedures
 analysis of variance, findings of
 age, 212-218, 301, 302
 behavior, 271-274
 employment state, 315-317
 etiological factors, 323
 institutionalized patients, 320-322
 IQ change
 Full Scale, 290-294
 Performance, 298-300
- Statistical procedures (continued)
 analysis of variance, findings of (con-
 tinued)
 remission, long-term, 207-212
 success of treatment in hospital,
 226t, 227t, 228-233, 230t, 231t,
 234-236, 235t, 236t
 discriminant function analysis for
 predicting
 behavior, 274-276
 intellectual deterioration, 302-306
 remission, long-term, 254-257
 seizure control in hospital, 237-
 239, 241-254
 cross validations, 239-240, 254-
 261
 factor analysis, 189-192, 203-207,
 266-267, 280
 intercorrelation, 185-189, 264-266,
 267-271, 278-280, 281-288, 314,
 315
 t test for nonindependent means, IQ
 change, 289
- Status epilepticus, 28, 33, 98, 190t,
 196, 201, 324
 age, 216
 as cause of death, 157-162, 165-170,
 194t
 remission of seizures, 11, 28, 191,
 201t, 202, 208, 211t, 228
- Steroids
 ACTH, treatment of infantile spasms,
 77, 78, 80-84
 corticosteroids, 82-83
 corticotrophin, 79, 80, 81
 Cortisone, 80, 81
 Prednisolone, 79, 80, 81
 Prednisone, 81
- Stigmata of degeneration
 mental deterioration, 126, 147
- Succinimides, 63
- Suicide, 160, 163, 166, 170
- Surgical treatment, 106-125
 age at onset of illness, 109, 121
 barbiturate fast activity, 121
 brain damage, diffuse, 121
 cerebral atrophy, bilateral, 121
 complications, 119

Surgical treatment (continued)
 electroencephalogram
 dysrhythmia, generalized, 121
 focus(i)
 unilateral versus bilateral, 120
 postoperative prognosis, 120, 121
 sphenoidal spike focus, 121
 focal epilepsy
 review of concept underlying surgery for, 115
 follow-up
 criteria used at, 115, 116, 118
 length of, 118
 improvement of mentality after, 107
 mesial temporal sclerosis, 121
 mortality, 108, 109, 113, 115, 119
 overt local changes in the brain, 109, 121
 personality, 121
 pneumoencephalography, 121
 prognostic criteria, 121
 results of, 108, 109, 110-111t, 112, 113, 114t, 115, 116, 117t, 118, 118t, 120, 122, 123, 124
 type of procedure
 excision
 cortical, 108, 112
 frontal lobe, 122
 of scars, 108
 hemispherectomy, 124
 Kocher valve, 112
 Pallidoamygdalotomy, 123
 Pallidoansotomy, 123
 Pallidotomy, 123
 splitting of
 corpus callosum, 123
 dura, 108
 temporal lobectomy, 116-123, 117t, 118t, 149-150
 criteria of eligibility, 119
 long-term remission rate, 117, 118
 number of patients operated on, 119, 120

Surgical treatment (continued)
 type of procedure (continued)
 temporal lobotomy, 122
 trephination, 108, 112

T

Temporal lobectomy (see Surgical treatment)
 Temporal lobotomy (see Surgical treatment)
 Thinking slowness, 137
 Tonic seizures (see Seizures)
 Trance-like states, 39
 Treatment (see also Anticonvulsant and Surgical)
 results
 inpatient, Lafayette Clinic, 225-228
 outpatient, Lafayette Clinic, 220-223
 success of, in hospital
 analysis of variance, findings, 225-233, 234-237
 Tridione, 16

V

Valium, 86
 Vocational Rehabilitation Administration Project, 258-261, 317-318

W

Wechsler Test (see also IQ and Psychological tests)
 laterality of EEG focus and, 137, 153
 Weight discrimination tests (see also Psychological tests)
 EEG and, 152
 West syndrome, 74
 Work ability, 16, 122, 126 (see also Employment)